

ARTICLE

Received 5 Jun 2014 | Accepted 5 Jan 2015 | Published 6 Feb 2015

DOI: 10.1038/ncomms7202

The transcriptional regulator BBX24 impairs DELLA activity to promote shade avoidance in *Arabidopsis thaliana*

Carlos D. Crocco^{1,†}, Antonella Locascio^{2,3}, Cristian M. Escudero¹, David Alabadí², Miguel A. Blázquez² & Javier F. Botto¹

In response to canopy shade, plant vegetative structures elongate to gain access to light. However, the mechanism that allows a plastic transcriptional response to canopy shade light is not fully elucidated. Here we propose that the activity of PIF4, a key transcription factor in the shade signalling network, is modulated by the interplay between the BBX24 transcriptional regulator and DELLA proteins, which are negative regulators of the gibberellin (GA) signalling pathway. We show that GA-related targets are enriched among genes responsive to BBX24 under shade and that the shade-response defect in *bbx24* mutants is rescued by a GA treatment that promotes DELLA degradation. BBX24 physically interacts with DELLA proteins and alleviates DELLA-mediated repression of PIF4 activity. The proposed molecular mechanism provides reversible regulation of the activity of a key transcription factor that may prove especially relevant under fluctuating light conditions.

¹IFEVA, Facultad de Agronomía, Universidad de Buenos Aires y Consejo Nacional de Investigaciones Científicas y Técnicas, Avenue San Martín 4453, Buenos Aires C1417DSE, Argentina. ²Instituto de Biología Molecular y Celular de Plantas, Consejo Superior de Investigaciones Científicas-Universidad Politécnica de Valencia, Valencia 46022, Spain. ³Department of Agronomy, Food, Natural Resources, Animals and the Environment (DAFNAE), Padova University, Viale dell'Università 16, Legnaro 35020, Italy. † Present address: Department of Botany and Plant Biology, University of Geneva, Sciences III, 30 Quai E. Ansermet, Geneva 4, 1211, Switzerland. Correspondence and requests for materials should be addressed to J.F.B. (email: botto@agro.uba.ar).

Light is a highly heterogeneous environmental factor controlling plant growth and survival, and defines the competitive relationships within the plant community¹. Contrasting light gradients occur in plant canopies, so all plants are exposed to at least some degree of shade during their lifetime. Sun plants have evolved complex signalling mechanisms to avoid shade through the promotion of developmental responses known as the shade-avoidance syndrome (SAS) that includes, among other responses, hypocotyl and petiole elongation, the upward orientation of leaves and the reduction of branching². The relevance and the plasticity of this adaptive response are illustrated by the huge genetic variation that is found in nature^{3–6}.

The canopy light is rich in far-red (FR) and poor in red (R) and blue light. Shade-avoiding species have developed a precise machinery to finely perceive changes in R:FR ratio through the R/FR-light-absorbing phytochrome family of photoreceptors². In open-light environments, when the R:FR ratio is high, the red-absorbing cytosolic form of phytochrome B, (phyB; Pr, max. absorbance = 660 nm) is excited and photoconverted to a FR-absorbing form (Pfr, max. absorbance = 730 nm) that moves to the nucleus, where it physically interacts with basic helix-loop-helix (bHLH) transcription factors called PHYTOCHROME-INTERACTING FACTORS (PIFs). Rapid phosphorylation and degradation of PIFs induces a dramatic reduction of growth rate in hypocotyls, stems and petioles⁷. In contrast, in shaded environments, the R:FR ratio is low and consequently the phytochromes are converted to the Pr form with low affinity for PIFs and preferential accumulation in the cytosol, allowing these transcription factors to accumulate and bind the promoters of cell elongation genes⁸.

Transcriptional regulation in response to shade is not only regulated by PIFs. In fact, a small network of transcription factors, including the non-DNA-binding bHLHs LONG HYPOCOTYL IN FR LIGHT (HFR1), PHYTOCHROME RAPIDLY REGULATED1 (PAR1) and PAR2 modulate this process preventing an exaggerated shade-avoidance response by forming inactive heterodimers with PIFs^{9–11}. Furthermore, double B-Box (BBX) containing zinc-finger transcription factors perform contrasting roles in the shade-avoidance response¹⁰. For instance, BBX21/STH2 and BBX22/STH3 act as negative regulators, while BBX24/STO and BBX25/STH1 promote hypocotyl growth of *Arabidopsis* seedlings under shade^{11,12}. Moreover, all BBX members studied so far are involved in the CONSTITUTIVE PHOTOMORPHOGENIC1 signalling pathway promoting the shade response^{11,12}.

Several hormones are also essential components in the response to shade, owing to their role as central players in the regulation of cell elongation. Auxin, gibberellins (GAs) and brassinosteroids promote elongation². For instance, a low R:FR ratio enhances both GA biosynthesis and responsiveness in *Arabidopsis*, thereby promoting the expression of GA-related genes^{13,14}. GA signalling proceeds through the promotion of the degradation of DELLA proteins, a small family of nuclear-localized transcriptional regulators that repress GA-responses, such as cell elongation. The molecular mechanism by which DELLAs regulate transcription is through the physical interaction with transcription factors, including PIFs¹⁵. In open environments, DELLAs accumulate and inactivate PIF4, thus preventing cell elongation^{16,17}. In agreement with this, enhanced petiole elongation under shade is accompanied by DELLA degradation¹⁸. Under low R:FR, GA biosynthesis is enhanced by the upregulation of *GA20ox* and *GA3ox* expression in petioles of *A. thaliana*¹⁴ and *Rumex palustris*¹⁹. However, despite the function of GA-regulating cell elongation responses is well established, the GA signalling pathway operating under shade is uncertain²⁰.

Despite the importance of BBX proteins in the SAS¹⁰, the molecular mechanism by which BBX proteins exert their action under shade also remains to be determined. Here, we demonstrate that BBX24 is involved in the GA-branch promoting cell elongation under shade. In particular, our results show that BBX24 physically interacts with, and inactivates, DELLA proteins, thereby providing a flexible mechanism that explains the positive role of BBX24 in the control of shade-elongation responses.

Results

BBX24 regulates hormone-related genes in response to shade.

To determine the relative importance of BBX24 in the SAS, we examined the hypocotyl elongation when *bbx24* and *BBX24*-overexpressing (*BBX24ox*) seedlings were grown under white light (high R:FR) and simulated shade (low R:FR). Two *bbx24* mutant alleles conferred normal hypocotyl elongation under high R:FR, but this ability was significantly reduced under low R:FR (Fig. 1). In addition, hypocotyls were longer in *BBX24ox* seedlings than in the wild type under high and low R:FR, but still responded to simulated shade (Fig. 1). These results demonstrate that BBX24 promotes hypocotyl elongation, but also other molecular elements are needed for a full response to shade. In fact, BBX25, its homologue, is one of these additional elements acting in parallel to BBX24. Although the *bbx25-2* allele did not show defects in response to low R:FR, the *bbx24-1 bbx25-2* double mutant displayed a greater reduction of the elongation response than each of the single mutants (Fig. 1). Furthermore, a simulated canopy treatment that reduces both the irradiation and R/FR significantly altered the *bbx25* seedling phenotype that failed to fully respond (Supplementary Fig. 1). Again, the *bbx25* allele was able to enhance the impaired phenotype of *bbx24* in the double mutant (Supplementary Fig. 1).

To identify genes regulated by BBX24 under simulated shade, we compared the transcriptomes of wild-type and *bbx24-1* seedlings exposed to high or low R:FR. We set a false-discovery rate of 4.12% and 1.5-fold change as cutoff to select genes putatively regulated by BBX24. Among a total of 432 genes, 49 were misregulated independently of the light quality (that is, genotype effect), 84 genes were differentially expressed in the *bbx24* mutant under high R:FR, and 299 genes were significantly altered under low R:FR (shade-regulated genes; Fig. 2a and Supplementary Data 1). The group of shade-regulated genes was




Figure 1 | Hypocotyl length in wild-type (Col-O and Ws), *bbx24* and *bbx25* single and double mutants and *BBX24ox* seedlings grown under white light or simulated shade (high and low R:FR ratios, respectively) in a chamber for 5 days. Bars indicate mean \pm sem ($n = 20$). **indicate differences between indicated means with $P \leq 0.01$ by Student's *t* test.




Figure 2 | BBX24 regulates 299 genes under shade. (a) Venn diagram showing the number of genes regulated by BBX24 under white light and shade. (b) Functional classification of the 299 genes regulated by BBX24 under shade. A bootstrap analysis was done to estimate the over-representation of each functional class and error bars, and the *P* values are indicated. (c) Cluster analysis of BBX24-regulated genes under shade using the D-chip program: transcription factors, signalling and hormone metabolism categories are shown.

enriched in gene categories associated with stress responses, cell-wall activity, secondary metabolism, hormone metabolism and transport-activity functions (Fig. 2b). D-chip analysis shows that shade-regulated genes can be clustered by transcription factors, signalling and hormone elements (Fig. 2c). Given that BBX proteins have been reported to regulate transcription through the interaction with DNA-binding transcription factors^{12,21,22}, we searched for common regulatory motifs in the promoters of the shade-regulated genes that could indicate BBX24 partners in this process (Supplementary Data 2). Among 299 genes regulated by BBX24 under simulated shade, we found that 212 genes (71%) contained one or two GA-response elements (125 and 87, respectively), and 77 genes (26%) contained auxin-response elements. Additionally, I-Box, G-Box, CCA1 and SORLIP1 elements were enriched (24, 23, 19, and 15% respectively; Supplementary Data 2). The link between BBX24 and hormone target genes under shade is also supported by the over-representation of auxin signalling genes previously linked to SAS, such as *IAA29* and *SAUR68* (Supplementary Fig. S2 and Supplementary Data 1). Auxin conjugation (*GH3.3*), GA catabolism (*GA2ox1*, *GA2ox2*), ethylene synthesis (*ACS2*) and circadian clock (*ELF4*) genes were also regulated by BBX24 under shade (Supplementary Fig. 2). Interestingly, *GA20ox2* and *GA3ox1* genes that are required for the synthesis of active GAs do not belong to the group of BBX24 shade-regulated genes (Supplementary Data 2), but their expression in *bbx24* seedlings exposed to shade is significantly different to wild type by quantitative reverse-transcription-PCR (RT-qPCR) analysis (Supplementary Fig. 2). In summary, the global expression analysis suggests that BBX24 can act directly or indirectly in conjunction with hormonal and light/circadian clock networks to promote hypocotyl elongation under shade.

GAs restore elongation of *bbx24* hypocotyls under shade. Enhanced production of active auxins and GAs is required for

proper hypocotyl growth in response to low R:FR^{18,23}. As a significant proportion of genes regulated by BBX24 under low R:FR belong to the auxin and GA pathways (Fig. 2 and Supplementary Fig. 2, Supplementary Data 1 and 2), we examined the effects of hormone levels to evaluate which of them is responsible for the *bbx24* phenotype. Picloram, a synthetic auxin (0.5 or 1 μ M), was equally effective in promoting hypocotyl elongation in wild-type and in *bbx24* mutant seedlings (Fig. 3a and Supplementary Fig. 3). In a complementary manner, 1-N-naphthylphthalamic acid (NPA, 0.5 μ M), an inhibitor of polar-auxin transport, treatment reduced the elongation in wild-type and *bbx24* seedlings under high and low R:FR, indicating the shade response similar between both genotypes (Fig. 3a and Supplementary Fig. 4). Other hormones involved in promoting hypocotyl elongation in the SAS, such as ethylene and brassinosteroids^{23,24}, did not rescue the mutant phenotype of *bbx24-1* and *bbx24-2* seedlings under low R:FR (Supplementary Fig. 3).

Remarkably, we found that the defective hypocotyl elongation phenotype of *bbx24-1* and *bbx24-2* seedlings under shade was fully rescued by the addition of GA₃ (Fig. 3a). To further support the functional relationship between BBX24 and the GA pathway, we performed a GA₃ dose-curve response (Fig. 3b). *bbx24-1* and *bbx24-2* seedlings were hypersensitive to GA₃ and showed significant differences compared with the wild type at 1 μ M that further increased at 5 μ M. In contrast, *BBX24ox* seedlings showed less sensitivity to the exogenous application of GA₃ than the wild type. These results suggest that BBX24 is involved in the regulation of GA metabolism and/or signalling under shade.

BBX24 promotes shade avoidance through PIF4. PIF4 is a central component of the transcriptional network regulating growth and it integrates diverse signals, including light and GAs^{16,17}. Furthermore, PIF4 promotes SAS through binding to the G-box (CACGTG) in the promoter of shade genes⁹.




Figure 3 | Gibberellins (GAs) restore hypocotyl length of *bbx24* seedlings under shade. (a) GA₃, but not picloram (auxin), restores the hypocotyl length of *bbx24* mutant seedlings under shade. paclobutrazol (PAC; GA synthesis inhibitor), but not 1-N-naphthylphthalamic acid (NPA; auxin transport inhibitor), suppresses the hypocotyl elongation mediated by BBX24 under shade. (b) GA₃ dose-response curve for hypocotyl growth promotion relative to mock (%) in wild-type, two *bbx24* mutant alleles and *BBX24ox* transgenic lines. Bars in **a** and points in **b** indicate means \pm s.e.m. ($n = 20$). * $P \leq 0.05$ and ** $P \leq 0.01$ by Student's *t*-test. NS, not significant differences between indicated means.

Interestingly, promoters of BBX24-regulated genes are enriched in the same *cis*-element (Supplementary Data 2). Then, we hypothesized that BBX24 is implicated in the PIF4-dependent shade signalling. To have a better understanding of this possible functional connection we generated *bbx24 pif4* double mutant. Hypocotyls of *pif4*, *bbx24* and *pif4 bbx24* mutants were very similar to each other and shorter than the wild type under low R:FR (Fig. 4a). Furthermore, we compared the *Arabidopsis* transcriptomes regulated by BBX24 (our data), PIF4 and PIF7 when seedlings were grown under low R:FR^{25–27}. The meta-analysis of 299 BBX24-regulated and 246 PIF-regulated genes under shade found that 16 genes were co-regulated by BBX24 and

PIFs (Fig. 4b and Supplementary Data 3). We observed a significant and positive association between genes regulated by BBX24 and PIFs (representation factor = 4.8, $P < 2.87e - 0.7$). These results suggest that BBX24 and PIF transcription factors are involved in the same signalling pathway and act in the same direction promoting elongation. Furthermore, we compared BBX24 shade-regulated genes with the PIF4 targets obtained by chromatin immunoprecipitation²⁸. Remarkably, 27 genes (9%) are both bound *in vivo* by PIF4 and commonly regulated by both proteins (Supplementary Data 4). By RT-qPCR analysis, we confirmed that the expression of genes upregulated by BBX24 under shade, *IAA29*, *SAUR68* and *GH3.3*, are downregulated in *bbx24*, *pif4* and *bbx24 pif4* mutants in a similar manner. *PIL1*, a gene not represented in the ATH1 chip, but known to be an early shade-regulated gene, also showed a reduced expression in single and double mutants of *BBX24* and *PIF4* (Fig. 4c). Conversely, plants overexpressing *BBX24* showed an enhanced induction of *IAA29*, *SAUR68* and *GH3.3* gene expression in response to shade (Fig. 4c). The results of both expression and physiological experiments suggest that BBX24 and PIF4 participate in the same signalling pathway co-regulating a common group of genes that promote elongation under shade.

BBX24 interacts with DELLAs to relieve repression of PIF4. Previous evidence clearly demonstrate that much of the DELLAs' involvement in the control of cell expansion is exerted through the modulation of PIFs' activity^{16,17}. The observations that the addition of GA₃ in the growth medium recovers the *bbx24* phenotype and the large overlap between PIF4 and BBX24 transcriptomes indirectly link BBX24 function to DELLA activity. Indeed, we found that this functional connection was correlated with the physical interaction between BBX24 and GAI and RGA, the two major DELLAs controlling hypocotyl elongation (Fig. 5a)²⁹. The results of yeast-two-hybrid assays with truncated BBX24 protein did not identify any particular interacting domain, suggesting that BBX24 might establish several and independent contacts with M5-GAI (Fig. 5b). A similar analysis with deleted versions of GAI highlighted the importance of the leucine heptad repeat 1 (Fig. 5b) for the interaction, as reported for other partners^{16,17,30}. Importantly, we also confirmed the interaction in plant cells. YFN-BBX24 and YFC-GAI interacted in nuclei of epidermal cells of *Nicotiana benthamiana* leaves as shown by means of bimolecular fluorescence complementation (BiFC) assays, whereas fluorescence in nuclei of control leaves co-expressing YFN-BBX24/YFC and YFN/YFC-GAI pairs was below detection limits (Fig. 5c). In addition, we confirmed this physical interaction by co-immunoprecipitation of BBX24-YFP and GAI-RFP transiently co-expressed in *N. benthamiana* leaves (Fig. 4d, Supplementary Fig. 5).

At least two possible mechanistic models based on the interaction between DELLAs and BBX24 could explain the positive role of BBX24 in the SAS: either BBX24 is a negative regulator of DELLA activity, thereby alleviating the repression of PIF4 by DELLAs, or alternatively, BBX24 is a co-activator of PIF4 that is negatively regulated by DELLAs. To distinguish between these two hypotheses, we set up a transient expression assay in *N. benthamiana* leaves using *PIL1::LUC* as a reporter for the activity of PIF4 (ref. 31), given that *PIL1* is a direct target of transcriptional regulation by PIF4 (ref. 32), and importantly, is also regulated by BBX24 activity under shade (Fig. 4c). As expected, expression of PIF4 increased *PIL1::LUC* activity at least threefold with respect to basal levels, and this induction was strongly reverted by co-expression of GAI (Fig. 6a). Interestingly, PIF4 activity was not altered by co-expression of BBX24, ruling




Figure 4 | BBX24 promotes shade avoidance through PIF4. (a) Hypocotyl length of wild-type, *bbx24* and *pif4* single and double mutant seedlings grown under white light and simulated shade. (b) Co-regulated genes by BBX24 (data of this work) and PIFs under shade (1) Li *et al.*²⁵, (2) Leivar *et al.*²⁷, (3) Hornitschek *et al.*²⁶ (c) *IAA29*, *SAUR68*, *GH3.3* and *PIL1* expression measured by RT-qPCR showing co-regulation by BBX24 and PIF4 under shade. The transcript levels were standardized to the wild type under white light. Bars in **a** and **c** indicate means \pm s.e.m. ($n = 20$ and $n = 3$, respectively). $***P \leq 0.001$ by Student's *t*-test. NS, not significant differences between indicated means.




Figure 5 | BBX24 interacts physically with DELLAs. (a, b) Yeast two-hybrid (Y2H) assay analysing the interaction between BBX24 and GAI and RGA. (c) Bimolecular Fluorescence Complementation (BiFC) analysis in *N. benthamiana* leaves of BBX24 and GAI fusions to N- and C-terminal fragments of YFP, respectively (upper photographs). Nuclei of control leaves co-expressing YFN-BBX24/YFC and YFN/YFC-GAI pairs are shown in the bottom photographs. Scale bars, 40 and 16 μ m (d) Co-immunoprecipitation assay showing the interaction between BBX24 and GAI in leaves of *N. benthamiana*.




Figure 6 | BBX24-GAI protein interaction promotes PIF4 activity.

(a) Transient expression assay in *N. benthamiana* leaves using *PIL1::LUC* as a reporter ('cps' are the Firefly Luciferase (LUC) fluorescence 'counts per sample', relative to the control Renilla Luciferase fluorescence used to normalize agro infiltration. BBX24 expression alleviates the repression of PIF4 by GAI. Bars indicate means \pm s.e.m. ($n = 2$ or 3). $*P \leq 0.05$ by Student's *t*-test, NS, not significant. (b) Bimolecular Fluorescence Complementation (BiFC) assay between PIF4 and GAI in *N. benthamiana* leaves is reverted by co-expression of BBX24. The bottom panels show that BBX24, GAI and PIF4 are expressed even if no BiFC signal is detected, according to the immunodetection of HA-BBX24 and semiquantitative PCR. Scale bars, 40 μ m.

out that this protein acts as a direct co-regulator of PIF4 activity. However, co-expression of BBX24 alleviated the repression imposed by GAI upon PIF4 (Fig. 6a), suggesting that this could be the mechanism by which BBX24 promotes the transcriptional response to shade through PIF4. Indeed, competition for DELLAs by PIF4 and BBX24 was further supported by BiFC experiments in which the interaction between PIF4 and GAI was impaired by co-expression of BBX24 (Fig. 6b).

To better understand the timing of BBX24-DELLA module action under shade, we measured the expression of some genes and DELLA protein levels in short-term shade. Seedlings of wild-type, *bbx24* and *BBX24ox* were cultivated in white light for 5 days and then exposed between 0 and 2 h to supplemental FR at the end of the photoperiod. We found that *PIL1*, *GH3.3* and *SAUR68* were dramatically and rapidly upregulated by simulated shade in wild-type seedlings and the absence of BBX24 significantly reduced their expression (Fig. 7a). Furthermore, RGA protein levels were not altered in wild-type, *bbx24* and *BBX24ox* seedlings cultivated for 2 h of short-term shade (Fig. 7b, Supplementary Fig. 5). As predicted by our BBX24-DELLA interaction model, these results suggest that the expression of BBX24-regulated genes under shade may be altered rapidly without the need for degradation of RGA proteins, a process that requires more time for the action of proteasome. However, the transcript levels of *RGA* and *PIF4* in short-term shade were not affected by the presence of BBX24 (Supplementary Fig. 6), we found that long-term shade significantly increased the expression of *DELLAs*, *PIF4* and *PIL1* mediated by BBX24 (Supplementary Fig. 7).

Our model also predicts that enhanced accumulation of DELLA proteins should cause resistance to the overexpression of BBX24 under shade conditions. To test this possibility, we examined the phenotype of *pGAI::gai-1D:GR* (*gai-1:GR*) seedlings expressing a conditional allele of GAI resistant to GA-induced degradation fused to the rat glucocorticoid receptor (GR) under the control of the native *GAI* promoter²⁸. The *gai-1:GR* fusion




Figure 7 | Expression of shade-regulated genes and DELLA protein levels in short-term simulated shade.

Seedlings of wild type, *bbx24* and *BBX24ox* were cultivated in white light for 5 days and then exposed between 0 and 2 h to supplemental far-red (FR) at the end of the photoperiod. (a) Shade-regulated genes change dramatically in short-term shade. *GH3.3*, *SAUR68* and *PIL1* expression were measured by RT-qPCR at 0, 0.5, 1 and 2 h after exposing seedlings to simulated shade. The transcript levels were standardized to the wild type under white light. Means \pm s.e.m. ($n = 4$). (b) RGA protein levels do not change in short-term shade in a western blot using anti-RGA antibody (1:300). C, control at $t = 0$ h; WL, white light at $t = 2$ h; +FR, simulated shade at $t = 2$ h.

protein is retained in the cytoplasm in the absence of the added steroid dexamethasone (DEX), whereas application of DEX results in its relocation from the cytoplasm to the nucleus. In the absence of DEX, hypocotyls of *gai-1* and *gai-1:GR* seedlings were shorter and similar to the wild-type seedlings under simulated shade, respectively (Fig. 8); similarly, the *gai-1:GR* transgene did not alter the phenotype of *BBX24ox* seedlings under the same light conditions (Fig. 8). As expected, DEX treatment caused an inhibition of hypocotyl growth under shade of *gai-1:GR* seedlings, which was then comparable with that of the *gai-1* mutant; and this inhibition was even more evident in the




Figure 8 | Hypocotyl length of wild-type, *gai-1*, *gai-1:GR*, *BBX24ox* and *gai-1:GR BBX24ox* mutant seedlings grown under white light and simulated shade in response to application of the steroid dexamethasone (+ DEX) or mock solution (- DEX). Bars indicate mean \pm s.e.m. ($n = 15$). *** $P \leq 0.001$ by Student's t -test. NS, not significant differences between indicated means.

BBX24ox background (Fig. 8), indicating that the promotion of elongation by *BBX24* under shade was in fact compromised by increasing DELLA's levels. This result is therefore in agreement with DELLAs acting downstream of *BBX24* in the shade response.

Discussion

An outstanding question in modern biology is how plants integrate environmental information in a plastic manner to modulate developmental habits. The evidence presented in this work provides a molecular mechanism that modulates the pathway linking the perception of shaded environments and the transcriptional regulation that ultimately results in a change in the growth pattern (Fig. 9). The mechanism by which *BBX24* regulates DELLA's activity in the context of shade avoidance is physiologically relevant, as it illustrates how light quality signals may relay the information through protein-protein interaction originating a small transcriptional regulatory network with well-defined flexible properties allowing a rapid and reversible response to shade. We have provided clear evidence that *bbx24* phenotype is rescued by GAs and operates in the GA signalling pathway by interrupting GAI-PIF4 interaction. In fact, we found that the impaired elongation phenotype of *bbx24* mutants is rescued by the addition of GA, and also the *bbx24* mutants are more sensitive to GA (Fig. 3). Nevertheless, the expression of some metabolic GA genes, targets of DELLA regulation, did not explain the phenotype of *bbx24* mutants. While *GA20ox2* and also *GA2ox2* were downregulated *GA3ox1* and *GA2ox1* were




Figure 9 | Model of *BBX24* action in shade signalling. In open environments with a high R:FR, the major form of phyB is PfrB, which destabilizes PIF4 and prevents the expression of shade-induced genes. In crowded stands with a low R:FR, the Pr form of phyB (PrB) increases and PIF4 binds to the G-box (CACGTG) in the promoter of genes that promote elongation. The PIF4 stability increases under shade because (i) PrB has low affinity to bind with PIF4 for degradation, (ii) PrB is mainly re-located to the cytosol, and (iii) *BBX24* decreases the amount of DELLAs capable to interact with PIF4. We speculate that an early or intermittent response to shade favors protein-protein reversible interactions between DELLAs and *BBX24*, while a prolonged shade favors quantitatively the GA-induced irreversible degradation of DELLAs by the proteasome machinery. The thickness of the arrow indicates the reaction favoured under shade.

upregulated by shade in the *bbx24* mutant (Supplementary Fig. 2). These results suggest that the feedback regulation of GA metabolic genes by DELLAs involves unknown elements also affected by *BBX24*. In addition, our results provide a novel case in which DELLA activity is downregulated by a mechanism different to GA-induced protein degradation mediated by the GID1 receptors as was demonstrated before³³. The results suggest that some DELLA repressors could be sequestered by *BBX24* proteins which would allow more PIF transcription factors to activate cell growth to elongate vegetative structures avoiding light competition under shade.

Owing to their activity as negative regulators of an increasing number of DNA-binding transcription factors, DELLAs act as general growth repressors that need to be inactivated to promote cell expansion^{15,34}, although it has also been shown that DELLAs can act as cofactors promoting the activity of some transcription factors³⁵⁻³⁷. Interestingly, all known environmental signals that promote growth induce the degradation of DELLA proteins by the proteasome. This is the case of higher ambient temperatures³⁸, darkness periods^{39,40} and also low R:FR ratios¹⁸, all of which presumably alter GA levels or the accumulation of GID1 receptors to drive to the reduction of DELLAs in the nucleus. However, the interaction of DELLA proteins with *BBX24* represents an alternative mechanism for the modulation of DELLA activity that does not involve protein degradation, but allows the sequestration of DELLA protein in competition with the PIF4 transcription factor. This regulatory cascade based on protein-protein interactions ultimately regulating gene expression represents an emerging theme in plant signalling pathways. For instance, a triple HLH/bHLH cascade has been described by which the DNA-binding ability of a bHLH transcription factors (PIF4 or ACE1) is negatively regulated by the physical interaction with an HLH protein (HFR1, PAR1 or IBH1), which in turn is repressed by direct interaction with another HLH protein such as PRE1 (refs. 41,42). An equivalent regulatory module has been reported for MYC2, whose transcriptional activation capacity is repressed by

interaction with the JAZ regulators, while DELLA proteins interact physically with JAZs to revert this repression^{30,43}. BBX proteins have been previously defined as transcriptional regulators^{12,21,22} with opposite regulating functions in shade and other developmental processes¹⁰. For example, BBX21 inhibits *ABI5* expression by interfering with HY5, and also, ABI5 protein binding to the *ABI5* gene promoter in germinating seeds²². Moreover, BBX24 and BBX25 interact with HY5 preventing its DNA-binding to the promoter of *BBX22* gene to induce its expression in seedling de-etiolation¹². In addition, BBX32 may interact physically with other BBX DNA-binding protein, like BBX21, and avoid the promotion of photomorphogenic genes²¹. In this context, our results show that BBX24 indirectly regulates the DNA-binding activity of PIF4 by the direct interference with DELLAs to promote cell elongation under shade.

An additional property of the BBX24-DELLA-PIF4 regulatory module, particularly important in the context of shade avoidance, is the intrinsic reversibility conferred by protein–protein interactions, compared with cascades involving *de novo* protein synthesis, or protein degradation. This plasticity may become critical in short-term or transient exposure to shade. In fact, the capacity of perception and transduction of light gap signals in a dynamic canopy environment to adjust the growth of vegetative structures is relevant to define the competitive relationships between plant individuals. Albeit the importance of kinetics in the shade growth response in different species, the knowledge of molecular mechanisms involved in each phase of growth elongation is scarce³⁰ and limited to the upregulation of some genes like *PIL1*, *ATHB2* or *HFR1* in short-term shade^{11,44}. Here, we propose a possible mechanism for a reversible and a rapid response to shade that involves the protein–protein interaction between BBX24 and DELLAs that competes for the interaction between DELLAs and PIF4.

One of the most important features inherent to the topology of the BBX24-DELLA-PIF4 module is that the transcriptional output of the cascade depends on the fine balance between molecular elements, which is determined by different and sometimes antagonistic signals that become integrated through protein–protein interactions. Given that DELLA and PIF4 levels are differentially influenced by temperature among other signals, and BBX24 and PIF4 stability has also been found to be dependent on light intensity, this regulatory module would render responses to shade in variable magnitudes depending on other environmental conditions and developmental phases. In fact, the shade-avoidance mechanisms may operate in different development processes including, between others, the control of seed germination, the orientation of branches and leaves, the elongation of stems, the production of new branches or tillers and the performance of the photosynthetic apparatus. Then, the capacity of plants to exploit rapid and efficiently the light opportunities in competition with neighbours would be a major determinant of evolutionary success. So it would be interesting to investigate if the molecular mechanism depicted here has been subject to adaptive pressure in different developmental situations and habitats.

Methods

Plant material. The mutants and transgenic lines used in this study have been described previously: *bbx24-1*, *bbx25-2*, *bbx24-1 bbx25-2* and *BBX24ox* in Col background, and *bbx24-2* in WS background^{12,45}; *gai-1:GR* lines⁴⁶, and the *PIF4ox* line¹⁷. *pi4-101* was obtained from the *Arabidopsis* Biological Resource Center. To generate *HA-BBX24* fusion, the pENTR-BBX24 was recombined by LR reaction (<http://www.invitrogen.com>) with pEarleyGate-204 (ref. 47). *gai-1:GR BBX24ox* lines were generated by crossing *HA-BBX24* with *gai-1:GR* lines and selected for homozygous progeny by PCR using primers listed in Supplementary Table 1.

Growth conditions and light treatments. In general we used the same experimental protocols described in Crocco *et al.*¹¹ Seeds were sown in clear plastic boxes on 0.8% agar/water, and incubated in darkness at 4 °C to reduce dormancy and homogenize germination. After 4 days, imbibed seeds were exposed to a red pulse and incubated in darkness for 24 h at 25 °C to induce germination. Then, the boxes with seedlings were transferred to white light for de-etiolation during 2 days. Three-day-old seedlings were exposed to white light or white light supplemented with FR lamps in a growth chamber at 22 °C settled with a short-day photoperiod (8 h light + 16 h dark) for 5 days. White light treatment consisted in mercury lamps (General Electric HR175/R/DX/FL39 mercury 33026) that established a high R:FR = 2.3. Simulated shade was generated with the addition of incandescent lamps (Philips, R19-100R20/FL/S) and two paolini filters with a red acetate (Paolini 2031; La Casa del Acetato, Buenos Aires, Argentina) placed laterally into the growth chamber that established a low R:FR = 0.35. The up radiation was similar for both light treatments (PAR = 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$). In addition, we used a simulated canopy condition that reduces both irradiance and R/FR ratio (PAR = 30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and R:FR = 0.30) by putting one paolini filter on the top of the boxes with seedlings. Spectral photon fluencies were estimated with a Li-Cor (Li-188B, LiCor Corp., Lincoln, NE, USA), and PAR and R:FR were measured using an SpectroSense2 attached with a SKR-1850SS2 light sensor (Skye Instruments Ltd., Powys, UK).

Hormone experiments. Seeds were sown in agar/water medium supplemented with synthetic hormones: indole-3-acetic acid, GA₃, precursor of ethylene (ACC) or brassinosteroids (EpiBL); or specific inhibitors: NPA and paclobutrazol (PAC) that inhibit the auxin polar transport and the endogenous GA synthesis, respectively. For DEX experiment, seeds were sown in agar/water on a filter paper. The experimental protocol was similar to those exposed before with little adjustments. To allow germination during PAC treatments, seeds were first sown and germinated for 24 h under white light on top of a wet filter paper, after which the filter paper was transferred to the MS plate containing PAC at the indicated concentrations. In the DEX experiments, 2 μM DEX (+DEX) or control solution with ethanol (–DEX) was added on the top of filter papers immediately before to start the light treatments.

RT-qPCR experiments. For RNA expression, 100 mg of fresh seedlings were harvested and frozen immediately in liquid nitrogen. Total RNA was extracted using an RNeasy plant mini kit (Qiagen, <http://www.qiagen.com>). Crude RNA preparations were treated with 10 units of RNase-free DNase I (<http://www.promega.com>), and the samples were purified according to the RNeasy plant mini kit protocol. cDNA was synthesized from 1.5 μg of DNA-free RNA template using an oligo(dT) primer and SuperScript II reverse transcriptase (<http://www.invitrogen.com>). RT-qPCR analysis was performed on an optical 96-well plate using SYBR Green PCR master mix (Applied Biosystems, <http://www.appliedbiosystems.com>) and an ABI PRISM 7500 real-time PCR system (<http://www.appliedbiosystems.com>). The thermal cycle used was 95 °C for 15 min, followed by 40 cycles of 95 °C for 15 s, 60 °C for 30 s and 72 °C for 35 s. Specific primer pairs for each gene were designed using Beacon Designer 7.0 (<http://www.premierbiosoft.com>) and are listed in Supplementary Table 1. *Actin8* (At1g49240), *UBC* (At5g25760) and *IPP2* (At3g02780) were used to normalize the expression levels and then standardized to the wild type under white light.

Microarrays and data analysis. The microarray experiment consisted in a factorial design comprising two genotypes (Col and *bbx24-1*) and two light treatments (white light and simulated shade). RNA material was extracted from 7-day-old seedlings. Two replicates per treatment were performed using Affymetrix *Arabidopsis* ATH1 GeneChip (<http://www.affymetrix.com>). RNA was prepared, labelled and hybridized in accordance with the manufacturer's instructions. Data were normalized by multiplying the value of each gene by the mean of each chip, divided by the average intensity of all mean values. Significantly differentially expressed genes were identified by performing profile analysis using Significance Analysis of Microarrays with a δ -value of 1.40, which corresponds to a false-discovery rate of 4.12% (ref. 48). Genes with 'absent' calls and a signal of < 50 units in all replicate experiments were filtered out. A test filter was performed to work only with those genes for which the ratio of expression showed at least a 1.5-fold change between Col and *bbx24-1*. Clusters were generated using DNA-Chip Analyzer (<http://www.dchip.org>). Gene Ontology enrichment were analysed using the Classification SuperViewer Tool of the Bio-Array Resource (<http://bar.utoronto.ca>). A ranking score is calculated for each functional class. The input set is bootstrapped 100 times to provide some idea as to over- or under-representation reliability. Upstream promoter sequences (1,000 bp) of differentially expressed genes were analysed using Promoter at BAR (<http://bar.utoronto.ca>). The *P* value was determined using the hypergeometric probability distribution to find the likelihood of the observed number of elements occurring in a randomly chosen set of promoters ($P < 0.05$). Statistical significance of the overlap between two groups of genes was calculated as the representation factor, which is the number of overlapping genes divided by the expected number of overlapping genes estimated from two independent groups (<http://elegans.uky.edu/>).

Protein–protein interaction assays. A pENTR vector carrying the coding sequence (CDS) of *BBX24* was created by cloning *BBX24* CDS into pCR8-GW vector (Invitrogen). Deleted versions of *BBX24* were amplified by PCR and cloned into pCR8-GW/TOPO (<http://www.invitrogen.com>) to create pENTR vectors. Deletions of GAI are described in Gallego-Bartolomé *et al.*⁴⁹ The full-length *BBX24* CDS was cloned into both pDEST22 and pDEST32, whereas deletions were cloned into pDEST22. Final bait and prey constructs were used to co-transform the yeast strain AH109 (<http://www.clontech.com>). Yeasts were selected in SD/-Leu/-Trp/-His and with different amounts of 3-aminotriazol (<http://www.sigmaaldrich.com>) to test interactions. Primers used for cloning are listed in Supplementary Table 1.

For BiFC, GAI and *BBX24* CDSs were transferred into pMDC43-YFC and pMDC43-YFN vectors, respectively⁵⁰. To study the interference of *BBX24* in the interaction between GAI and PIF4, the GAI and PIF4 CDSs were transferred to pMDC43-YFC and pMDC43-YFN vectors, respectively, whereas that of *BBX24* was transferred to pEarleyGate-201 (ref. 47) to create a HA fusion. BiFC analysis was performed as described in Locascio *et al.*¹⁵. The oligonucleotides used as primers to detect GAI, PIF4 and *EF1- α* from *N. benthamiana* leaves by RT-sqPCR have been described^{38,51}. For co-immunoprecipitation assays in *Nicotiana benthamiana*, *BBX24* CDS was transferred as indicated above into pEarleyGate-104 (ref. 47) to create the YFP-*BBX24* fusion. RFP-GAI is described in Locascio *et al.*¹⁵ Each construct was introduced into *Agrobacterium tumefaciens* C58 cells. Three days after infiltration, leaves were collected and frozen in liquid nitrogen. Infiltrated tissues were ground and homogenized in two volumes of cold extraction buffer (50 mM Tris · HCl (pH 7.5), 50 mM NaCl, 1% (vol/vol) Nonidet P-40 and 1 × complete protease inhibitor mixture (<http://www.roche.com>)). Extracts were centrifuged twice for 30 min at 16,000 × g in a top bench microcentrifuge at 4 °C. Total soluble proteins in the supernatant were quantified by Bradford's assay. Ten micrograms of soluble proteins were saved to be used as input, and the remaining proteins were incubated with 50 μ l of anti-GFP paramagnetic MicroBeads (<https://www.miltenyibiotec.com>) on ice for 2 h in a total volume of 1 ml. Extracts were loaded at room temperature onto μ -columns (<https://www.miltenyibiotec.com>) previously equilibrated with extraction buffer. Columns were kept at room temperature and washed four times with 200 μ l of cold extraction buffer and once with 100 μ l of cold washing buffer 2 (20 mM Tris · HCl, pH 7.5) supplied in the kit (<https://www.miltenyibiotec.com>). Proteins were eluted in 50 μ l of denaturing elution buffer following the manufacturer's instructions. Immunoprecipitated proteins were run in 12% SDS-polyacrylamide gel electrophoresis, immunoblotted and detected with anti-GAI antibody diluted 1:1,000 (ref. 52). Subsequently, blots were stripped out and incubated with anti-GFP antibody diluted 1:1,000 (Ab290; <https://www.abcam.com>). The western blot in Fig. 7b was incubated with an anti-RGA antibody diluted 1:300 (<http://www.agrisera.com>).

Reporter constructs and transcriptional assays. *pPIL1::LUC* reporter constructs are describe in Zhang *et al.*³¹ As effector proteins, we used PIF4-YFP, *BBX24*-HA and GAI-GFP fusions¹⁵. To generate PIF4-YFP fusion, the pENTR carrying PIF4 CDS from REGIA (Regulatory Gene Initiative in *Arabidopsis*)⁵³ was recombined by LR reaction into a pEarleyGate-104 (ref. 47). *BBX24*-HA construct was made by recombining the pENTR carrying *BBX24* into the vector pEarleyGate-201 by the LR reaction. Transient expression in leaves of 4-week-old *N. benthamiana* was done by the infiltration mixture. To prevent silencing, *A. tumefaciens* C58 carrying a construct that expresses the silencing suppressor P19 was included in the mixtures. The ratio of cells carrying P19:reporter:effector constructs was 1:1:1 for PIF4 and *BBX24*, whereas it was 1:1:4 for GAI. Mixtures were incubated for 3 h in darkness at RT before infiltration. Firefly and the control Renilla-LUC activities were assayed from leaf extracts collected 3 days after infiltration with the Dual-Glo Luciferase Assay System (Promega) and quantified with a GloMax 96 Microplate Luminometer (Promega).

References

- Valladares, F. & Niinemets, U. Shade tolerance, a key plant feature of complex nature and consequences. *Annu. Rev. Ecol. Evol. Syst.* **39**, 237–257 (2008).
- Casal, J. J. Photoreceptor signaling networks in plant responses to shade. *Annu. Rev. Plant Biol.* **64**, 403–427 (2013).
- Botto, J. F. & Coluccio, M. P. Seasonal and plant-density dependency for quantitative trait loci affecting flowering time in multiple populations of *Arabidopsis thaliana*. *Plant Cell Environ.* **30**, 1465–1479 (2007).
- Coluccio, M. P., Sánchez, S., Kasulin, L., Yanovsky, M. J. & Botto, J. F. Genetic mapping of natural variation in a shade avoidance response: ELF3 is the candidate gene for a QTL in hypocotyl growth regulation. *J. Exp. Bot.* **62**, 167–176 (2011).
- Filiault, D. L. & Maloof, J. N. A genome-wide association study identifies variants underlying the *Arabidopsis thaliana* shade avoidance response. *PLoS. Genet.* **8**, e1002589 (2012).
- Kasulin, L., Agrofoglio, Y. & Botto, J. F. The receptor-like kinase ERECTA contributes to the shade-avoidance syndrome in a background-dependent manner. *Ann. Bot.* **111**, 811–819 (2013).
- Leivar, P. & Monte, E. PIFs: systems integrators in plant development. *Plant Cell* **26**, 56–78 (2014).
- Lorrain, S., Allen, T., Duek, P. D., Whitelam, G. C. & Fankhauser, C. Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant J.* **53**, 312–323 (2008).
- Hornitschek, P., Lorrain, S., Zoete, V., Michielin, O. & Fankhauser, C. Inhibition of the shade avoidance response by formation of non-DNA binding bHLH heterodimers. *EMBO J.* **28**, 3893–3902 (2009).
- Gangappa, S. N. & Botto, J. F. The BBX family of plant transcription factors. *Trends Plant Sci.* **19**, 460–470 (2014).
- Crocco, C. D., Holm, M., Yanovsky, M. J. & Botto, J. F. AtBBX21 and COP1 genetically interact in the regulation of shade avoidance. *Plant J.* **64**, 551–562 (2010).
- Gangappa, S. N. *et al.* The Arabidopsis B-BOX protein BBX25 interacts with HY5, negatively regulating BBX22 expression to suppress seedling photomorphogenesis. *Plant Cell* **25**, 1243–1257 (2013).
- Devlin, F. P., Yanovsky, M. J. & Kay, S. A. A genomic analysis of the shade avoidance response in *Arabidopsis*. *Plant Physiol.* **133**, 1–13 (2003).
- Hisamatsu, T., King, R. W., Helliwell, C. A. & Koshioka, M. The involvement of gibberellin 20-oxidase genes in phytochrome-regulated petiole elongation of *Arabidopsis*. *Plant Physiol.* **138**, 1106–1116 (2005).
- Locascio, A., Blázquez, M. A. & Alabadi, D. Genomic analysis of DELLA protein activity. *Plant Cell Physiol.* **54**, 1229–1237 (2013).
- de Lucas, M. *et al.* A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**, 480–486 (2008).
- Feng, S. *et al.* Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**, 475–480 (2008).
- Djakovic-Petrovic, T., de Wit, M., Voeselek, L. A. C. J. & Pierik, R. DELLA protein function in growth responses to canopy signals. *Plant J.* **51**, 117–126 (2007).
- Pierik, R., de Wit, M. & Voeselek, L. A. C. J. Growth-mediated stress escape: convergence of signal transduction pathways activated upon exposure to two different environmental stresses. *New Phytol.* **189**, 122–134 (2011).
- Colebrook, E. H., Thomas, S. G., Phillips, A. L. & Hedden, P. The role of gibberellin signalling in plant responses to abiotic stress. *J. Exp. Biol.* **217**, 67–75 (2014).
- Holtan, H. E. *et al.* BBX32, an Arabidopsis B-Box protein, functions in light signaling by suppressing HY5-regulated gene expression and interacting with STH2/BBX21. *Plant Physiol.* **156**, 2109–2123 (2011).
- Xu, D. *et al.* Convergence of light and ABA signaling on the *ABI5* promoter. *PLoS. Genet.* **10**, e1004197 (2014).
- Pierik, R., Djakovic-Petrovic, T., Keuskamp, D. H., de Wit, M. & Voeselek, L. A. C. J. Auxin and ethylene regulate elongation responses to neighbor proximity signals independent of gibberellin and DELLA proteins in *Arabidopsis*. *Plant Physiol.* **149**, 1701–1712 (2009).
- Keuskamp, D. H. *et al.* Blue-light-mediated shade avoidance requires combined auxin and brassinosteroid action in *Arabidopsis* seedlings. *Plant J.* **67**, 208–217 (2011).
- Li, L. *et al.* Linking photoreceptor excitation to changes in plant architecture. *Genes Dev.* **26**, 785–790 (2012).
- Hornitschek, P. *et al.* Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling. *Plant J.* **71**, 699–711 (2012).
- Leivar, P. *et al.* Dynamic antagonism between phytochromes and PIF family basic helix-loop-helix factors induces selective reciprocal responses to light and shade in a rapidly responsive transcriptional network in *Arabidopsis*. *Plant Cell* **24**, 1398–1419 (2012).
- Oh, E., Zhu, J.-Y. & Wang, Z.-Y. Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nat. Cell Biol.* **14**, 802–809 (2012).
- Dill, A. & Sun, T. P. Synergistic derepression of gibberellin signaling by removing RGA and GAI function in *Arabidopsis thaliana*. *Genetics* **159**, 777–785 (2001).
- Cole, B., Kay, S. A. & Chory, J. Automated analysis of hypocotyl growth dynamics during shade avoidance in *Arabidopsis*. *Plant J.* **65**, 991–1000 (2011).
- Zhang, Y. *et al.* A quartet of PIF bHLH factors provides a transcriptionally centered signaling hub that regulates seedling morphogenesis through differential expression-patterning of shared target genes in *Arabidopsis*. *PLoS. Genet.* **9**, e1003244 (2013).
- Leivar, P. *et al.* Definition of early transcriptional circuitry involved in light-induced reversal of PIF-imposed repression of photomorphogenesis in young *Arabidopsis* seedlings. *Plant Cell* **21**, 3535–3553 (2009).
- Willige, B. C. *et al.* The DELLA domain of GA INSENSITIVE mediates the interaction with the GA INSENSITIVE DWARF1A gibberellin receptor of *Arabidopsis*. *Plant Cell* **19**, 1209–1220 (2007).
- Davière, J.-M. & Achard, P. Gibberellin signaling in plants. *Develop* **140**, 1147–1151 (2013).

35. Lim, S. *et al.* ABA-INSENSITIVE3, ABA-INSENSITIVE5, and DELLAs interact to activate the expression of SOMNUS and other high-temperature-inducible genes in imbibed seeds in *Arabidopsis*. *Plant Cell* **25**, 4863–4878 (2013).
36. Yoshida, H. *et al.* DELLA protein functions as a transcriptional activator through the DNA binding of the indeterminate domain family proteins. *Proc. Natl Acad. Sci. USA* **111**, 7861–7866 (2014).
37. Yamaguchi, N. *et al.* Gibberellin acts positively then negatively to control onset of flower formation in *Arabidopsis*. *Science* **344**, 638–641 (2014).
38. Stavang, J. *et al.* Hormonal regulation of temperature-induced growth in *Arabidopsis*. *Plant J.* **60**, 589–601 (2009).
39. Achard, P. *et al.* DELLAs contribute to plant photomorphogenesis. *Plant Physiol.* **143**, 1163–1172 (2007).
40. Arana, M. V., Marín-de la Rosa, N., Maloof, J. N., Blázquez, M. A. & Alabadi, D. Circadian oscillation of gibberellin signaling in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **108**, 9292–9297 (2011).
41. Bai, M.-Y., Fan, M., Oh, E. & Wang, Z.-Y. A triple helix-loop-helix/basic helix-loop-helix cascade controls cell elongation downstream of multiple hormonal and environmental signaling pathways in *Arabidopsis*. *Plant Cell* **24**, 4917–4929 (2012).
42. Ikeda, M., Fujiwara, S., Mitsuda, N. & Ohme-Takagi, M. A triantagonistic basic helix-loop-helix system regulates cell elongation in *Arabidopsis*. *Plant Cell* **24**, 4483–4497 (2012).
43. Yang, D.-L. *et al.* Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade. *Proc. Natl Acad. Sci. USA* **109**, E1192–E1200 (2012).
44. Ciolfi, A. *et al.* Dynamics of the shade-avoidance response in *Arabidopsis*. *Plant Physiol.* **163**, 331–353 (2013).
45. Indorf, M., Cordero, J., Neuhaus, G. & Rodríguez-Franco, M. Salt tolerance (STO), a stress-related protein, has a major role in light signalling. *Plant J.* **51**, 563–574 (2007).
46. Gallego-Bartolomé, J., Kami, C., Fankhauser, C., Alabadi, D. & Blázquez, M. A. A hormonal regulatory module that provides flexibility to tropic responses. *Plant Physiol.* **156**, 1819–1825 (2011).
47. Earley, K. W. *et al.* Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J.* **45**, 616–629 (2006).
48. Tusher, V. G., Tibshirani, R. & Chu, G. Significance analysis of microarrays applied to the ionizing radiation response. *Proc. Natl Acad. Sci. USA* **98**, 5116–5121 (2001).
49. Gallego-Bartolomé, J. *et al.* Molecular mechanism for the interaction between gibberellin and brassinosteroid signaling pathways in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **109**, 13446–13451 (2012).
50. Belda-Palazón, B. *et al.* Aminopropyltransferases involved in polyamine biosynthesis localize preferentially in the nucleus of plant cells. *PLoS ONE* **7**, e46907 (2012).
51. Gallego-Bartolomé, J., Alabadi, D. & Blázquez, M. A. DELLA-induced early transcriptional changes during etiolated development in *Arabidopsis thaliana*. *PLoS ONE* **6**, e23918 (2011).
52. Piskurewicz, U. *et al.* The gibberellic acid signaling repressor RGL2 inhibits *Arabidopsis* seed germination by stimulating abscisic acid synthesis and ABI5 activity. *Plant Cell* **20**, 2729–2745 (2008).
53. Paz-Ares, J. REGIA, an EU project on functional genomics of transcription factors from *Arabidopsis thaliana*. *Comp. Funct. Genomics* **3**, 102–108 (2002).

Acknowledgements

We thank Santiago Mora García for valuable initial discussions and Peter Quail for the *PL1::LUC* construct. This work was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica, and Universidad de Buenos Aires (to J.F.B.), and the Spanish Ministry of Science, BIO2010-15071 (to M.A.B.).

Author contributions

C.D.C., D.A., M.A.B. and J.F.B. designed research, C.D.C., C.E. and A.L. performed research; C.D.C., A.L., D.A., M.A.B. and J.F.B. analysed data; C.D.C., D.A., M.A.B. and J.F.B. wrote the paper.

Additional information

Accession codes: microarray data associated with this study has been deposited in the NCBI GEO database under the accession code GSE 64755.

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Crocco, C. D. *et al.* The transcriptional regulator BBX24 impairs DELLA activity to promote shade avoidance in *Arabidopsis thaliana*. *Nat. Commun.* **6**:6202 doi: 10.1038/ncomms7202 (2015).