

Long-term and recent changes in temperature-based agroclimatic indices in Argentina

María Elena Fernández-Long,^{a,*} Gabriela Viviana Müller,^b Adriana Beltrán-Przekurat^c
and Olga Eugenia Scarpati^{d,e}

^a Department of Agricultural Climatology, School of Agronomy, University of Buenos Aires, Buenos Aires, Argentina

^b Center of Scientific Research and Technological Transfer to Productivity/National Research Council (CICYTTP-CONICET), Diamante, Entre Ríos, Argentina

^c Department of Atmospheric and Oceanic Sciences and Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

^d Centro de Estudios Farmacológicos y Botánicos (Pharmacological and Botanical Center)/National Research Council (CEFYO-CONICET), Buenos Aires, Argentina

^e Geography Department, La Plata National University, La Plata, Argentina

ABSTRACT: Changes in several temperature-based agroclimatic indices in the central-eastern of Argentina, most of them located within the Pampas region were analysed for 39 meteorological stations. Trends of first (FFD), last (LFD) and number (NFD) of frost days, frost period (FP), start (SGS), end (EGS) and length (LGS) of the growing season, growing degree days (GDD), diurnal temperature range (DTR), chilling hours and lowest annual minimum temperature were computed for two periods, 1940–2007 and 1975–2007. The largest changes were observed for the whole period 1940–2007 and were mostly indicative of a long-term minimum temperature warming throughout the region. During this period, generalized decreases in the NFD and in the FP (i.e., a delayed FFD and an earlier LFD) were found. Although the trends in the growing season indices were not as large as in the frost indices, they were consistent with the overall warming: an earlier SGS and a delayed in the EGS. The trends of the GDD showed a large variability between months with a generalized increased throughout the year. The DTR showed the largest number of stations with statistically significant negative trends from austral late spring (November) to austral early fall (April). For the period 1975–2007, the behaviour changes in all analysed indices: the short-term trends weakened and in some cases reversed sign. The LFD tended to occurred later in the year, particularly for the southern Pampas. The EGS shifted from mostly positive to negative trends, resulting in a shorter LGS. These trend changes were not spatially homogeneous. Although those short-term trends were predominantly non-statistically significant, they could potential affect management decisions and crop yields. In particular, frost is still an important hazard in agricultural activities and within the context of our results short- and long-term characterization of frost risk need to be considered at local and sub-regional scales. Copyright © 2012 Royal Meteorological Society

KEY WORDS agroclimatic indices; climate change; Pampas region; temperature

Received 30 December 2010; Revised 30 May 2012; Accepted 3 June 2012

1. Introduction

Climate change has gained significant international attention over the past decade due to concerns of deleterious long-term impacts on agriculture, water supply, human welfare and regional political stability. Climate change is expected to affect extremes, since small changes in mean conditions can lead to large changes in the frequency of extremes (Katz and Brown, 1992). The length of the frost-free season has increased in most mid- and high-latitude regions of both hemispheres. In the Northern Hemisphere, this is mostly manifest as an earlier start to spring (Trenberth *et al.*, 2007).

Agricultural production and related activities (e.g., irrigation, fertilization, site selection) can be highly affected by long-term changes in temperature and precipitation. Agrometeorological indices can be used to evaluate how weather and climate conditions affect plant status and crop production (Alexandrov *et al.*, 2008), and can also help in evaluating management decisions and farm technologies (Eitzinger *et al.*, 2009).

Agrometeorological indices also have been used to evaluate changes in climate conditions. For example, long-term changes of temperature-based agrometeorological indices, like frost days and length of the growing season (GS), have been documented around the world. An overall decrease in the number of frost days has been found in different regions. Frich *et al.* (2002) found evidence of fewer frost days in most of mid- and high-latitudes in the Northern Hemisphere in the last 50 years. A decreasing number of frost days were found across

*Correspondence to: M. E. Fernández-Long, Department of Agricultural Climatology, School of Agronomy, University of Buenos Aires, Av. San Martín 4453, C1417DSE, Buenos Aires, Argentina. E-mail: flong@agro.uba.ar

northern Europe during the 20th century (Heino *et al.*, 1999), in Canada (Bonsal *et al.*, 2001) and in New Zealand (Salinger and Griffiths, 2001). In China, the last spring frost day has been occurring earlier since the 1980s (Yaodong, 2005). Nevertheless, for the southeast of United States, Easterling (2002) showed no changes in either the number of frost days or the frost-free season, in spite of a cooling trend in annual temperatures over the 20th century in that region.

Different regions around the world have shown a lengthening of the GS, related to an increase in the near-surface air temperature (Myneni *et al.*, 1997; Tucker *et al.*, 2001; Moonen *et al.*, 2002; Walther *et al.*, 2002; Root *et al.*, 2003; Shen *et al.*, 2005; Schwartz *et al.*, 2006). Walther and Linderholm (2006) showed that in the Greater Baltic Area for the 20th century the general mean trend of the length of the GS is about 20 d. During 1951–2000, the frost-free period has increased around 0.5 d per year in Germany, Austria and Switzerland and 0.34 d per year in Estonia (Menzel *et al.*, 2003). Feng and Hu (2004) showed that the thermal time significantly increased at annual rate of up to 70 degree days per 10 years across the western United States but decreased 20 degree days per 10 years from the US Great Plains to the east coast.

Changes in the diurnal temperature range (DTR), another agrometeorological index, have been linked to changes in crop yields. Lobell (2007) analyzed the historical yield data of rice and maize from the leading global producers and showed that an increase in DTR was associated with reduced yields. Increased DTR for a given average temperature may reduce yields because the associated increase in maximum temperature may result in a higher water stress or reductions in photosynthesis rates (Dhakhwa and Campbell, 1998). Similarly, reductions in minimum temperature associated with increased DTR may also be harmful in cases where freezing temperatures can result in crop injury or death (Rosenzweig and Tubiello, 1996; Tubiello *et al.*, 2002). For South America, Vincent *et al.* (2005) found a generalized decreased in the DTR over the continent mostly due to nighttime warming for the period 1961–2000.

This work is focused on the temperate region of South America, specifically the eastern part of Argentina (Figure 1). Most of the stations analysed in this work are located within the Pampas which represents the main agricultural region of Argentina and one of the major agricultural regions in the world (Hall *et al.*, 1992). Two additional stations outside the Pampas were added to the analysis. Those stations are situated in the north and northwestern part of Argentina, areas that experienced an exponential increase in the cultivated soya bean area and production (Grau *et al.*, 2005; Boletta *et al.*, 2006): the average soya bean production increased from 300 10³ ton in the late 1990s to 2.5 10⁶ ton in recent years (SIIA, 2010).

During the last two decades crop yields have significantly increased, greatly influenced by changes in




Figure 1. Studied region and meteorological stations used in this work (see Table I for station names).

technology and regional climate. Some of the technology improvements include: use of higher yielding and disease resistant varieties, and boost up of fertilization practices. In addition, increase in soil water availability was an essential contributor to the observed crop yield increases (Hurtado, 2008; Spescha, 2008). In the studied region, 38, 18 and 13% of the simulated yields increases for soya bean, maize and wheat between the periods 1950–1970 and 1971–1999 were associated with changes in climate (Magrin *et al.*, 2005). Within the context of future climate change, modelling results showed that increases in temperature could lead to decreases of potential yields of wheat, maize and sunflower (Magrin *et al.*, 1998) which may be offset, at least to a limited extent, by an increase in atmospheric carbon dioxide (CO₂) concentration (Magrin and Travasso, 2002).

Large-scale atmospheric circulation changes have occurred in this area, superimposed with a range of land-use land-cover changes that have affected temperature (Minetti and Vargas, 1983, 1992; Hoffmann *et al.*, 1997; Rusticucci and Barrucand, 2004; Fernández-Long and Müller, 2006; Nuñez *et al.*, 2008) and precipitation (Barros *et al.*, 2000; Rusticucci and Penalba, 2000; Minetti *et al.*, 2003; Haylock *et al.*, 2006; Forte Lay *et al.*, 2008). Rusticucci and Barrucand (2004) studied the changes in temperature extremes in Argentina over the period 1959–1998. They found that the largest and positive trends occurred in the summer and for the minimum temperatures. Particularly over the central-eastern Argentina,

the Pampas region, Fernández-Long and Müller (2006) examined annual and monthly trends in the number of frost days from 1964 to 2003. They found a decrease, although not statistically significant, in the annual frost frequency, and a large spatial variability in the trends.

Extreme temperatures can have a great impact on the crop production. In the case of summer crops, like soya bean maize or sunflower, their sowing date is conditioned upon the date of the last frost (Raper and Kramer, 1987; Otegui *et al.*, 1996; Otegui and Lopez Pereira, 2003). In the case of winter crops, like wheat, delays in the last frost can greatly affect the yields (Abbate *et al.*, 1994; Otegui and Lopez Pereira, 2003). In recent years, the occurrence of last frosts in November greatly affected the wheat production in the southern Pampas (Quiroz and Maneiro, 2007; Ross *et al.*, 2007). In addition, there was a generalized decrease in corn yields and some areas with soya bean had to be re-sown due to frost damage (Downes, 2007).

A preliminary analysis of the daily temperatures, i.e. maximum (T_{\max}) and minimum (T_{\min}) for selected stations within the studied area (Table I) shows generalized negative and positive long-term trends for the period 1940–2007 (Figure 2(a) and (b)). In addition, for the recent period 1975–2007 the T_{\max} and T_{\min} trends show a change in sign or a decreased in value. The number of stations with statistically significant negative T_{\min} tendencies decreased from 31 for the whole period 1940–2007 to only 11 stations for the shorter 1975–2007 period. Moreover, for the latter period, four stations showed statistically significant positive trends. A large spatial heterogeneity of this behaviour is found within the region. For example, for a station located south of the area (Figure 2(c) and (d)) T_{\max} trend change in sign, and T_{\min} trend change in absolute value becoming statistically significant. However, for a western station the short-term trends remained unchanged or changed slightly (Figure 2(e) and (f)).

Several questions arise given the potential effects of changes in daily extreme temperatures in the agricultural production:

1. What are the long-term changes of the main temperature-based agrometeorological indices in this region?
2. How those agrometeorological indices have changed in the recent years?
3. What is the spatial variability of those changes?
4. Are the changes related to global climate change?

The objective of this work is to evaluate the temporal and spatial changes of the main temperature-based agrometeorological indices in the eastern region of Argentina. We analysed the trends in the indices for two periods: 1940–2007 and 1975–2007. The latter was chosen considering our preliminary analysis of the T_{\max} and T_{\min} trends and the observed changes in atmospheric circulation from mid-1970s in different regions of the world (Trenberth and Hurrell, 1994; Nakamura *et al.*,

1997; Wang *et al.*, 2007, 2010) and in Argentina (Rusticucci and Penalba, 2000). In this paper Section 2 describes the data, indices and methodology; the trend analysis is presented in Section 3, and discussion and conclusions follow in Section 4.

2. Observational data and methodology

2.1. Climatic data

Daily minimum (T_{\min}) and maximum (T_{\max}) temperature from 39 stations in Argentina (Table I, Figure 1) with the longest and complete records were used. In addition these stations have little urban heat island effect and they were not relocated.

Two selected periods of data were used. The first one includes almost the total period of data of all considered meteorological stations. Only three stations allowed for the trend analysis for more than 80 years (Table I). Therefore, in order to include as many stations as possible, the period 1940–2007 was analysed. A second period, 1975–2007, was also considered based on the observed changes in atmospheric circulation around mid-1970s.

2.2. Agrometeorological indices

Table II lists the definitions of the agrometeorological indices analyzed in this study. In the case of frost-based indices, the first (FFD) and last (LFD) frost day, the frost period (FP) and the number of frost (NFD) days, we used the 0 and 3 °C thresholds. The FFD is defined as the first date on or before 15 July on which T_{\min} is less or equal than the temperature threshold. The LFD is the last date on or after 16 July on which T_{\min} is less or equal than the temperature threshold. The FP is the number of days between the FFD and LFD and NFD is the number of days with frost.

The length of the growing season (LGS) is another widely used index. In most mid-latitude regions, temperature is often the sole parameter to determine LGS, and in general, the 5 °C mean temperature (T_{mean}) threshold is widely accepted for determining the thermal growing season, in particular for mid- and high-latitudes (Jones and Briffa, 1995; Carter 1998; Frich *et al.*, 2002; Jones *et al.*, 2002; Menzel *et al.*, 2003). There are different LGS definitions, based on the length of the spells this threshold has to be exceeded or fall below. In our study, LGS is defined as the period from the start (SGS) to the end (EGS) of the growing season. The SGS is specified to be the first appearance of seven consecutive days with the T_{mean} higher than 5, 10 or 15 °C after the last winter/spring frost (Jones *et al.*, 2002). The EGS is defined to be the last appearance of seven consecutive days with T_{mean} higher than 5, 10 or 15 °C before the first autumn/winter frost. As elaborated in Peterson and Folland (2000), this definition of LGS is particularly relevant for measuring change of agricultural environment. The indices using the 5 °C base temperature coincided mostly with the frost-free period and therefore were not analysed.




Figure 2. Annual average (standardized) of daily maximum (left panels) and minimum temperature (right panels) and their trends for the periods 1940–2007 and 1975–2007: (a) and (b) for all the stations; (c) and (d) for Coronel Suarez (CSu) and (e) and (f) for Santa Rosa (SRo).

Crops grow when the daily T_{mean} is above a given temperature threshold, varying according to the specie and its phenological state. Different indices are used to quantify that process. Thermal time or growing degree day (GDD) has been used to assist in selections of crops and hybrids to assure the selected crops will achieve maximum growth at the time they reach maturity and potential yield. Relationships between crop growth and GDD also have been used to predict crop yield (Mall *et al.*, 2000; León *et al.*, 2001). Changes in monthly GDD could affect the duration of some phenological stages, and in particular the ones that could decisively affect crop production. Therefore, we decided to analyse the monthly GDD values. GDD for each month and for three different

temperature thresholds, or base temperature T_b , 5, 10 and 15 °C, GDD_{T_b} , were computed according to:

$$\text{GDD}_{T_b} = \sum_1^n (T_{\text{mean}} - T_b) \quad (1)$$

where n is the number of days in a given month, and T_{mean} is the daily average temperature computed using daily T_{min} and T_{max} . There is no accumulation if $T_{\text{mean}} < T_{\text{base}}$. We are not including in the analysis the GDD for base temperatures over 15 °C for the winter months (June to August).

Most plants from temperate regions, including deciduous fruit trees, require accumulating a given amount of

Table I. Meteorological stations (abbreviation and name), latitude (Lat), longitude (Lon), elevation (m) and initial year of available data.

Abbreviation	Station	Lat (S)	Lon (W)	Elevation (m)	Year
Sal	Salta	24.85	65.48	1221	1925
SdE	Santiago del Estero	27.77	64.30	199	1931
BVi	Bella Vista	28.43	58.92	70	1928
Cer	Ceres	29.88	61.95	88	1931
VMRS	Villa María del Río Seco	29.90	63.68	341	1931
MCa	Monte Caseros	30.27	57.65	54	1931
CorA	Córdoba	31.32	64.22	474	1956
Pil	Pilar	31.67	63.88	338	1925
Par A	Paraná	31.78	60.48	78	1931
VDo	Villa Dolores	31.95	65.13	569	1930
Ros	Rosario	32.92	60.78	25	1935
Gua	Gualeguaychú	33.00	58.62	21	1931
RCu	Río Cuarto	33.12	64.23	421	1931
SLu	San Luis	33.27	66.35	713	1931
Per	Pergamino	33.93	60.55	65	1931
Lab	Laboulaye	34.13	63.37	137	1939
SMi	San Miguel	34.55	58.73	26	1933
Eze	Ezeiza	34.82	58.53	20	1956
LPl	La Plata	34.97	57.90	23	1956
PIn	Punta Indio	35.37	57.28	22	1956
NdJ	Nueve de Julio	35.45	60.88	76	1931
Dol	Dolores	36.35	57.73	9	1931
SRo	Santa Rosa	36.57	64.27	191	1937
CSu	Coronel Suarez	37.43	61.88	233	1936
Pig	Pigüé	37.60	62.38	304	1935
Bal	Balcarce	37.75	58.30	130	1961
MdP	Mar del Plata	37.93	57.58	21	1950
BBl	Bahía Blanca	38.73	62.17	83	1956
Rec	Reconquista	29.18	59.70	53	1948
Tan	Tandil	37.23	59.25	175	1960
GPi	General Pico	35.70	63.75	141	1956
Cas	Castelar	34.67	58.65	22	1959
MJu	Marcos Juarez	32.70	62.15	114	1952
HAs	Hilario Ascasubi	39.38	62.62	22	1961
SVi	Sauce Viejo	31.70	60.82	18	1958
Jun	Junín	34.55	60.92	81	1958
Peh	Pehuajó	35.87	61.90	87	1951
Con	Concordia	31.30	58.02	38	1963
TAr	Tres Arroyos	38.33	60.25	109	1964

Table II. Agrometeorological indices used.

Indices	Abbreviation	Definition	Unit
First frost day	FFD	Date of first autumn frost	Date
Last frost day	LFD	Date of last spring frost	Date
Frost period	FP	Number of days between FFD and LDF	Days
Number of frost days	NFD	Number of days below freezing	Days
Start of growing season	SGS _{5,10,15}	First 7 d spell with T_{mean} above 5, 10 and 15 °C	Date
End of growing season	EGS _{5,10,15}	Last 7 d spell with T_{means} above 5, 10 and 15 °C	Date
Length of growing season	LGS _{5,10,15}	Number of days between SGS _{5,10,15} and EGS _{5,10,15}	Days
Growing degree days (monthly)	GDD _{5,10,15}	Accumulated number for temperature degrees above 5, 10 and 15 °C	°C
Diurnal temperature range (monthly)	DTR	Monthly difference between daily maximum and minimum temperature	°C
Chilling hours	CH	Number of hours below 7 °C	Hours
Lowest minimum temperature	LMT	Lowest minimum temperature of the year	°C

chilling temperatures to overcome a dormancy period and to resume the growth in spring. A very common index is the chilling hours (CH) defined as the total number of hours when the temperature is below or equal to 7 °C (Fraisse and Whidden, 2010). The daily CH were accumulated over the five coldest months, considering here from May to September, when the accumulation of chilling temperatures is maximum. CH from daily temperature data was estimated according to Pascale and Damario (2004):

$$\text{CH} = 24 \times \frac{(7 - T_{\min})}{(T_{\max} - T_{\min})} \quad (2)$$

The DTR for each month was calculated as the monthly average of the daily thermal amplitude, i.e. daily difference between T_{\max} and T_{\min} . The last index in the analysis is the lowest minimum temperature of the year (LMT) which represents an indication of the risk of lethal temperatures on a given crop (Hall and Sadras, 2009).

2.3. Data analysis

The least squares estimator of a regression coefficient is vulnerable to gross errors and the associated confidence interval is, in addition, sensitive to non-normality of the parent distribution (Sen, 1968). So, in this paper we calculate the presence of a monotonic increasing or decreasing trend in the agrometeorological indices with the nonparametric Mann–Kendall test (Mann, 1945; Kendall, 1975) and the slope of a linear trend is estimated with the nonparametric Sen's method (Sen, 1968). The trend is the median of all possible trend estimates obtained from a pairs of observations obtained at distinctly different observing times. This estimator is statistically robust and unbiased. Missing values are allowed and the data need not conform to any particular distribution. Besides, the Sen's method is not greatly affected by single data errors or outliers.

Zhang *et al.* (2004) confirm that the Mann–Kendall test and Sen's slope estimator are not reliable when applied on time series with serially correlated residuals. The time series analysed in this work have an annual time step; we considered them statistically independent and without autoregressive processes. The level of significance was set to $p < 0.05$. Hereafter, if no level of significance is specified, all trends significant at the level $p < 0.05$ are referred to as significant. The trends were computed for annual time series for all the indices and for each of the monthly time series for GDD and DTR.

3. Results

3.1. Frost indices

Period 1940–2007. The FFD over 0 °C showed a delay in most of the region: in 35 stations (90%) the trends were positive indicating that the FFD occurred later in the year (Table III, Figure 3(a)). In 14 stations (36%), the correlation was statistically significant. For these stations, the occurrence of the FFD was delayed on average 4 d

Table III. Number of stations with positive (Pos) and negative (Neg) trends of the frost indices FFD, LFD FP and NFD, above 0 and 3 °C, for the periods 1940–2007 and 1975–2007.^a

Period	Indices	Above 0 °C		Above 3 °C	
		Pos	Neg	Pos	Neg
1940–2007	FFD	35 (14)	4 (0)	35 (20)	4 (1)
	LFD	9 (0)	28 (13)	9 (0)	30 (18)
	FP	8 (0)	31 (16)	8 (1)	31 (23)
	NFD	9 (1)	30 (17)	4 (2)	35 (20)
1975–2007	FFD	25 (3)	14 (1)	23 (2)	16 (1)
	LFD	18 (4)	19 (1)	19 (1)	20 (1)
	FP	19 (2)	20 (4)	19 (3)	20 (4)
	NFD	20 (2)	19 (2)	16 (2)	23 (5)

^a Between parentheses is the number of stations with statistically significant trends.

decade⁻¹. On the other hand, the LFD in 28 stations (72%) showed negative trends (33% were statistically significant), indicating a generalized earlier occurrence of the last frost (Table III, Figure 3(c)). Most of the stations with non-statistically significant changes in the LFD index were located north of 30 °S and along 65 °W (Figure 3(c)). On average, the LFD trend statistically significant was -4 d decade⁻¹. The behaviour of both indices has resulted in a decrease of the FP, with a clear negative trend throughout the region, statistically significant in 16 stations (41%) (Table III, Figure 3(e)). The average of the statistically significant trends values was -7 d decade⁻¹.

There were also a lower number of frost days: the NFD decreased in 30 stations (77%), and in 17 (44%) the changes are statistically significant. The NFD statistically significant trends values ranged between -3 and 5 d decade⁻¹, with an average of -2 d decade⁻¹. In summary, from 1940 to 2007, fewer frost days occurred in most of the region. A small area in the SE of the studied region, centred approximately at 37 °S and between 62° and 59°W (stations CSu and Tan), shows the opposite behaviour, an increase number of frost days, within a longer time period (Figure 3(e) and (g)).

The trends of the frost indices over the 3 °C threshold were statistically significant in a larger number of stations than over the 0 °C threshold (Table III). The spatial variation is similar to the frost indices over 0 °C: an increase number of stations with statistically significant frost indices trends located SE part of the domain (not shown). The mean trend values were also similar to the values over the 0 °C threshold, but the range of variation was larger for the FFD and consequently for the FP. The FFD trends ranged between 2 and 5 d decade⁻¹ on average for the 0 °C threshold, compared to between -8 and 12 d decade⁻¹ for the 3 °C threshold. The FP trends ranged between -12 and -3 d decade⁻¹ for the 0 °C threshold and -16 and 12 d decade⁻¹ for the 3 °C threshold.

Period 1975–2007. All the indices in this shorter and recent period showed statistically significant trends in a




Figure 3. Trend of (a, b) FFD, (c, d) LFD, (e, f) FP and (g, h) NFD, above 0°C for the periods 1940–2007 (top) and 1975–2007 (bottom). Positive and negative trends are represented with circles and triangles, respectively. Stations with statistically significant trends are shown with closed dark symbols.

much lower number of stations (Table III, Figure 3, bottom). Moreover, the trends did not show a clear behaviour and even the spatial pattern changed throughout the region compared to the period 1940–2007 (Figure 3, top). For example, for the FFD index over 0°C , there were only 3 stations, compared to 14 in the longer period, with statistically significant positive trends. A 36% of the stations (14 stations) presented negative trends, compared to only 10% in the 1940–2007 period (Table III) and most of them located south of 35°S (Figure 3(b)). This means an earlier occurrence of the first frost during recent years. In general, the mean trend values are larger, in absolute values, than in the longer period 1940–2007. One station, Tandil (Tan), have now statistically significant negative trends. The LFD showed a similar change: 18 stations have now positive trends (9 in the longer period) and only one with a negative statistically significant trend, compared to 13 in the period 1940–2007 (Table III, Figure 3(d)).

Although the FP index over 0°C , showed a similar behaviour to the total period, the change was strongly weakened, in the slope value and in the number of stations with significant values (Table III, Figure 3(f)). The number of stations with statistically significant negative trends decreased from 16 to 4, and increased from none to 2 the stations with positive trends. Compared to the longer period (1940–2007), more stations show a relative longer FP during the recent years. This is indicated by the changes in the trends between the two periods (Table III): they show a shift from negative to positive

values or trends become less negative. Nevertheless, the trends tended to shorten mostly in stations located W and N of the domain (Figure 3(f)).

Finally, for the NFD index over 0°C , fewer stations have statistically significant negative trends (from 17 to 2 stations, Table III). More stations have now positive trends (20 compared to 9 in the period 1940–2007) mainly in the centre of the domain (Figure 3(h)). This indicates a tendency to more days with frost in the recent period.

Similar changes were found for the frost indices calculated over 3°C (Table III). For example, only 2 stations have statistically significant positive FFD trends compared to 20 in the longer period. The LFD index that clearly had a negative trend in 77% of the stations (30) during the period 1940–2007, shifted to positive trends in 19 stations (49%), although only 1 is statistically significant. The number of stations with statistically significant negative trends in the FP index drastically decreased (from 23 to 4), but increased the number of stations with positive trends, from 8 to 19, of which 3 are statistically significant. Similar to the NFD over 0°C , there are more stations with positive trends of the NFD over 3°C in the recent period 1975–2007.

3.2. Indices based on crop temperature requirements

3.2.1. Growing season indices (SGS, EGS and LGS)

Period 1940–2007. In general for both analysed periods there was a fewer number of stations with statistically

significant trends in the growing season indices compared to the frost indices (Table IV). Nevertheless, a clear behaviour can be seen for the base temperature of 10 and 15 °C. The SGS₁₀ and SGS₁₅ indices showed negative trends in 28 (72%) and 30 (77%) stations, although in only 12 and 9 the trends were statistically significant, respectively (Table IV). This indicates an earlier start of the growing season, on an average of -4.1 d decade⁻¹ for the SGS₁₀ index (only for the statistical significant stations). No station presented statistically significant positive trends (Table IV).

During this period, the trends in the EGS₁₀ and EGS₁₅ were positive in 33 and 30 stations, in 12 and 5 stations the trends were statistically significant (Table IV). This indicated a tendency to a delayed end of the growing season; the average value is 4 d decade⁻¹ for the EGS₁₀ index (only for the statistical significant stations). No station had negative statistically significant trends (Table IV).

The combination of both indices led to a general increase of the LGS throughout the region (Table IV). The LGS₁₀ and LGS₁₅ indices presented positive trends in 34 (87%) and 31 (79%) stations, and they were statistically significant in 18 and 14 stations, respectively. Although eight stations showed negative LGS₁₅ trends, none was statistically significant. For the LGS₁₀ the average increase is 6.3 d decade⁻¹ considering only the statistically significant stations, and 8.3 d decade⁻¹ for LGS₁₅. The trends did not have a clear spatial pattern in any of the indices (not shown).

Period 1975–2007. The number of stations with statistically significant trends decreased during this more recent period, similarly to the frost indices (Table IV). The SGS₁₀ and SGS₁₅ indices still showed negative trends, although number of stations with statistically significant trends shifted from 12 to none for SGS₁₀, and from 9 to 1 for SGS₁₅.

The EGS trends changed in this more recent period: most of the stations shifted from positive trends to negative trends, i.e., EGS₁₅ had 30 stations with positive

trends in 1940–2007, with 5 of them statistically significant, and in this more recent period 11 with only one statistically significant (Table IV). The trends values were still low, indicating that the end of the growing season tended to occur earlier.

The LGS₁₀ and LGS₁₅ indices shifted from predominantly positive trends during the period 1940–2007 (in 34 and 31 stations, respectively) to a lower number of stations with positive trends, (24 stations) for LGS₁₀, and mostly negative trends (24 stations) for LGS₁₅ in 1975–2007. The average LGS₁₀ value was -5.8 d decade⁻¹ for stations with negative trends. There were only two stations with statistically positive trends (Table IV). Similarly to the other growing season indices, the spatial distribution of the trends did not show a clear pattern, although the largest decrease in LGS tended to be concentrated in the centre of the domain (not shown).

3.2.2. Growing degree days

Period 1940–2007. The trends of the GDD index showed a large variability between months with a generalized increase throughout the year. The GDD₁₅ trends in January, March (Figure 4(a)), April and September to December were positive in more than 61% of the stations up to 97%, with no statistically significant negative trend, except in January (Table V). Except in March and October, the rest of the months had less than six stations with statistically significant positive trends. During October (Figure 4(e)), almost all the stations, 38 of 39, had positive trends, and 29 were statistically significant. The opposite pattern was observed in February and May (Figure 4(c)): they had mostly negative trends, with 28 and 33 stations respectively, and in February 7 were statistically significant (Table V).

The GDD₅ showed a similar behaviour than GDD₁₅. In June and August, most of the stations presented positive trends, with two and eight statistically significant (Table V). In July there were 34 stations with negative trends, although no station had statistically significant GDD₅ trend. October had all stations with positive trends and 32 were statistically significant. This was the month with the most spatially homogeneous trends.

Period 1975–2007. Similarly to the previously analysed indices, the number of stations with statistically significant GDD₁₅ and GDD₅ trends drastically decreased during this latest period (Table V). The monthly pattern of positive *versus* negative trends mostly remained the same compared to the period 1940–2007. A shift to more negative than positive trends, has occurred for the GDD₁₅ and GDD₅ in December. No statistically significant trends, either positive or negative, were found in the GDD₁₅ for March (Figure 4(b)), April and May (Figure 4(d)), and in the GDD₅ for March, May, July and August (Table V). The month with the largest number of stations with statistically significant trends was October (Figure 4(f)), although that number decreased from 29 and 32 for the GDD₁₅ and GDD₅ during the period 1940–2007 to 7 for this period 1975–2007 (Table V).

Table IV. Number of stations with positive (Pos) and negative (Neg) trends of the growing season indices SGS, EGS and LGS, above 10 and 15 °C, for the periods 1940–2007 and 1975–2007.^a

Period	Indices	Above 10 °C		Above 15 °C	
		Pos	Neg	Pos	Neg
1940–2007	SGS	10 (0)	28 (12)	9 (0)	30 (9)
	EGS	33 (12)	6 (0)	30 (5)	9 (0)
	LGS	34 (18)	5 (0)	31 (14)	8 (0)
1975–2007	SGS	18 (2)	20 (0)	18 (2)	21 (1)
	EGS	25 (4)	14 (2)	11 (1)	28 (2)
	LGS	24 (4)	15 (2)	15 (2)	24 (2)

^a Between parentheses is the number of stations with statistically significant trends.




Figure 4. Trend of GDD above 15 °C, for the periods 1940–2007 (top) and 1975–2007 (bottom) for (a, b) March, (c, d) July and (e, f) October. Positive and negative trends are represented with circles and triangles, respectively. Stations with statistically significant trends are shown with closed dark symbols.

Table V. Number of stations with positive (Pos) and negative (Neg) trends of GDD for the periods 1940–2007 and 1975–2007.^a

	Period 1940–2007				Period 1975–2007			
	GDD ₁₅		GDD ₅		GDD ₁₅		GDD ₅	
	Pos	Neg	Pos	Neg	Pos	Neg	Pos	Neg
January	26 (5)	13 (3)	25 (8)	14 (4)	20 (2)	19 (0)	20 (2)	19 (0)
February	11 (0)	28 (7)	12 (0)	27 (8)	10 (0)	29 (2)	10 (0)	29 (3)
March	36 (11)	3 (0)	35 (12)	4 (0)	33 (0)	6 (0)	32 (0)	7 (0)
April	30 (5)	9 (0)	30 (8)	9 (0)	23 (0)	16 (0)	19 (0)	20 (2)
May	6 (0)	33 (1)	11 (0)	28 (1)	3 (0)	36 (0)	3 (0)	36 (0)
June	–	–	32 (2)	7 (0)	–	–	37 (2)	2 (0)
July	–	–	5 (0)	34 (0)	–	–	14 (0)	25 (0)
August	–	–	38 (8)	1 (0)	–	–	30 (0)	9 (0)
September	30 (1)	9 (0)	30 (8)	9 (0)	28 (2)	11 (0)	30 (2)	9 (2)
October	38 (29)	1 (0)	39 (32)	0 (0)	38 (7)	1 (0)	35 (7)	4 (0)
November	26 (4)	13 (0)	26 (4)	13 (0)	36 (4)	3 (0)	31 (3)	8 (0)
December	24 (2)	15 (0)	25 (1)	14 (0)	12 (0)	27 (1)	12 (0)	27 (1)

^a Between parentheses is the number of stations with statistically significant trends.

February, with negative trends, also showed a similar behaviour in the recent years, but with less statistically significant trends.

3.3. Other agroclimatic indices

3.3.1. Diurnal temperature range

Period 1940–2007. The DTR presented the higher number of stations with statistically significant trends of all the analysed indices and clearly most of them are negative (Table VI, Figure 5). The DTR clearly decreased from austral spring (September to October to November) to austral autumn (March to April to May) over almost the whole domain. June and July had the lowest number of stations with statistically significant trends.

Period 1975–2007. The number of stations with statistically significant trends drastically decreased in this latest period (Table VI, Figure 5). Moreover, there is not a clear seasonal pattern as in the period 1940–2007. The number of months with statistically significant positive trends increased specially during in April to May (austral autumn) and August to September (austral late

Table VI. Number of stations with positive (Pos) and negative (Neg) trends of DTR for the periods 1940–2007 and 1975–2007.^a

	Period 1940–2007		Period 1975–2007	
	Pos	Neg	Pos	Neg
January	2 (0)	37 (32)	18 (1)	21 (1)
February	1 (0)	38 (34)	19 (0)	20 (1)
March	1 (0)	38 (23)	18 (0)	21 (1)
April	2 (0)	37 (27)	9 (2)	30 (7)
May	5 (1)	34 (13)	17 (2)	22 (4)
June	12 (2)	27 (2)	8 (1)	31 (2)
July	17 (4)	22 (4)	25 (3)	14 (0)
August	10 (0)	29 (9)	25 (5)	14 (0)
September	11 (0)	28 (7)	22 (3)	17 (0)
October	6 (1)	33 (12)	16 (0)	23 (0)
November	2 (0)	37 (22)	30 (1)	9 (0)
December	2 (0)	37 (25)	29 (1)	10 (0)

^a Between parentheses is the number of stations with statistically significant trends.




Figure 5. Trend of DTR for the periods 1940–2007 (top) and 1975–2007 (bottom) for (a, b) August, (c, d) October and (e, f) November. Positive and negative trends are represented with circles and triangles, respectively. Stations with statistically significant trends are shown with closed dark symbols.

winter – early spring). The pattern of trends started to shift to more positive than negative trends in August and extended to December, with the exception of October (Figure 5).

3.3.2. Chilling hours and lowest minimum temperature

Period 1940–2007. Negative trends in the CH were observed in 35 stations and in 11 were statistically significant (Table VII). Only four stations showed positive trends, and none statistically significant. This general decrease is due mostly to higher nighttime temperatures.

The LMT trends were positive in 30 stations, with 11 statistically significant (Table VII). Only two stations, CSu and Tan, showed negative statistically significant trends (not shown).

Period 1975–2007. Similarly to the rest of the indices, the number of stations with CH and LMT statistically significant trends decreased (Table VII). The pattern remained the same in both indices although the number of stations with CH negative trends decreased from 35 to 25 (Table VII).

4. Discussion and conclusions

In this work, we analysed the trends for the main temperature-based agroclimatic indices for two periods, 1940–2007 and 1975–2007 over the main agricultural area of Argentina. Overall, the trends changed between both studied periods, in absolute values and direction, although tendencies were mostly non-statistically significant in the short-term 1975–2007 period.

The occurrence of frost showed a clear decreasing trend in the long-term period 1940–2007 throughout the region. The FFD occurred later and the LFD occurred earlier during recent years compared to the 1940s and therefore, the frost period shortens during this total period 1940–2007. Conversely, a non-spatially homogeneous behaviour is observed during the recent 1975–2007 period. Most of the trends decreased. Moreover, in some stations located in the southern and central areas showed negative LFD trends, i.e. the last frost occurring later in the year. Although only few stations show statistically significant trends in this short-term period, the frost risk is still present with the consequent potential crop damage (Fernández-Long *et al.*, 2005; Magra *et al.*,

2005). During 2007 and 2008, frosts in November greatly affected the wheat production in the southern Pampas (Ross *et al.*, 2007; Quiroz and Maneiro, 2007) and some areas with soya bean had to be re-sown due to frost damage (Downes, 2007).

The long-term tendencies of the SGS and EGS were mostly negative and positive, respectively, indicating an earlier start and a delayed end of the growing season, respectively. Therefore, the LGS presented a positive trend for the two temperature thresholds used. These results are similar to the ones observed in other parts of the world, like in Germany (Menzel *et al.*, 2003), China (Song *et al.*, 2010), and in mid-latitudes in the Northern Hemisphere (Frich *et al.*, 2002). Increases of the LGS could also shift the productive area to higher latitudes, and also increase the number of potential harvest opportunities (ACIA, 2004). In addition, a longer LGS could add more flexibility to some agricultural practices which could lead to maximize yields (Rinaldi, 2004; Grassini *et al.*, 2011). The sowing date for summer crops is usually delayed due to high probabilities of occurrence of the last frost; therefore an earlier start of the growing season would allowed for an earlier sowing date of summer crops which would increase the possibility of double-cropping, for example, wheat–soya bean, as found for the south-eastern Pampas (Monzón *et al.*, 2007). They also found that a longer growing season, due to higher temperatures, resulted in increased simulated soya bean yield in the double-cropping management.

Nevertheless, during the recent 1975–2007 period, the EGS reverses the sign in many of the stations, and therefore, the growing season length shorten, which could have a negative impact on yield. Again, these short-term trends are statistically significant in few stations.

The GDD index trends showed a high inter-monthly variability. In the long-term period, an increase in temperature and therefore in GDD was found during the transitional months March and October (in more than 90% of the stations) while a temperature decrease was found during the extreme months July and February (in more than 69% of the stations). This means a seasonal change in the GDD index, with more availability of temperature in the sowing season and less for the reproductive season (February) of summer crops (i.e. maize and soya bean). More energy available during wheat flowering (October) could shorten the critical period and therefore potentially diminishing yields (Slafer and Rawson, 1994; Calderini *et al.*, 2001).

The GDD trends smoothed out during the recent period 1975–2007 compared to the total period and the number of stations with statistically significant trends drastically decreased in this latter period. GDD trends during austral summer months December and January SGS₁₀ and SGS₁₅ shifted from mostly positive to mostly negative values, indicating a decreasing of temperature accumulation during the first development stages of summer crops.

Similar behaviour was found for the DTR monthly index. In the long-term period, during austral summer,

Table VII. Number of stations with positive (Pos) and negative (Neg) trends of CH and LMT for the periods 1940–2007 and 1975–2007.^a

Period	Indices	Pos	Neg
1940–2007	CH	4 (0)	35 (11)
	LMT	30 (11)	9 (2)
1975–2007	CH	14 (0)	25 (1)
	LMT	31 (5)	8 (3)

^a Between parentheses is the number of stations with statistically significant trends.

in more than 64% of the stations the trends were negative and statistically significant. During austral winter, although a large number of stations showed negative trends, only a few were statistically significant. These results are similar to those found by Nuñez *et al.* (2008), Rusticucci and Barrucand (2004) and Vincent *et al.* (2005) and they can be explained by the increase in minimum temperature. The increase of T_{\min} (and the decrease of T_{\max}) can be explained by the increase of precipitation mainly during the summer semester (Forte Lay *et al.*, 2008). In the short-term period 1975–2007 there was a tendency to reverse the sign of the trends, especially during the warm spring-summer months, although similarly to the GDD index, the number of statistically significant trends drastically decreased. The annual precipitation trends showed a similar behaviour to the temperature-based indices, but of opposite value: all the analysed stations showed statistically significant positive values in the long-term annual precipitation tendency (Figure not shown). No station showed statistically significant precipitation tendencies for the short 1975–2007 period. A thorough analysis of the seasonal and monthly precipitation trends and their relationship with temperature trends is beyond the scope of the present work. We are deferring the analysis for a later study.

Instrumental observations over the past 157 years show that temperatures at the surface have risen globally, with important regional variations. For the global average, warming in the last century has occurred in two phases, from the 1910s to the 1940s (0.35 °C), and more strongly from the 1970s to the present (0.55 °C) according to the latest IPCC report (Trenberth *et al.*, 2007). Our results showed a different behaviour respect to the global increase in temperature: we found that the warming for the more recent 1975–2007 period was weaker than in the long-term 1940–2007 period, opposite to the global findings. For example, the LGS₁₅ trends in 31 of the 39 analysed stations were positive (in 14 stations the trends were statistically significant) during the long-term period; the sign of the trends reversed for the recent short-term period, and 24 stations had negative trend values (and only 2 were statistically significant).

The behaviour of the trends and the changes between the long- and short-term periods were mostly spatially homogeneous with a clear exception of the stations Tandil (Tan) and Coronel Suárez (CSu) located in the southern Pampas. They showed a decrease in the minimum temperature, especially during the short-term 1975–2007 period and a statistically significant increase of the frost period. This shows that changes at local and sub-regional scales can greatly differ from those of the overall region. The temporal evolution, long- and short-term, of agroclimatic indices and extreme temperatures, and their effects on agricultural production and management should be locally assessed.

In summary, long-term 1940–2007 trends of several temperature-based agroclimatic indices for eastern Argentina show a generalized indication of regional

warming, mainly due to increases in the minimum temperatures. For a recent period 1975–2007, the trends mostly weakened or reversed direction in some stations. Although those short-term trends are predominantly non-statistically significant, as Liebmann *et al.* (2010) found to be the case for short-term global surface temperature, they could potential affect management decisions and crop yields. In particular, frost is still an important hazard in agricultural activities and within the context of our results short- and long-term characterization of frost risk might need to be considered at local and sub-regional scales.

Acknowledgements

The meteorological information was provided by the Argentine National Weather Service and the Instituto Nacional de Tecnología Agropecuaria (INTA). This work was partially supported by grants CONICET (PIP-114-200801-00591). The authors wish to thank to anonymous reviewers whose insightful comments and suggestions led to improve the manuscript.

References

- Abbate PE, Andrade FH, Culot JP. 1994. Determinación del rendimiento de Trigo. Boletín Técnico 133, EEA INTA Balcarce. Bs. As., Argentina.
- ACIA. 2004. *Impacts of a Warming Arctic: Arctic Climate Impact Assessment*. Cambridge University Press: Cambridge, New York, 140.
- Alexandrov V, Mateescu E, Mestre A, Kepinska-Kasprzak M, Stefano VD, Dalezios N. 2008. Summarizing a questionnaire on trends of agroclimatic indices and simulation model outputs in Europe. Cost Action 734 Impact of Climate Change and Variability on European Agriculture – Survey of Agrometeorological Practices and Applications in Europe regarding Climate Change Impacts. P. Nejedlik and S. Orlandini, 115–161.
- Barros V, Castañeda ME, Doyle M. 2000. Recent precipitation trends in southern South America east of the Andes: an indication of climatic variability. In *Southern Hemisphere Paleo and Neoclimates*, Smolka PP, Volkheimer W (eds). Springer-Verlag: Berlin and Heidelberg, Germany, 187–206.
- Boletta P, Ravelo A, Planchuelo A, Grilli M. 2006. Assessing deforestation in the Argentine Chaco. *Forest Ecology and Management* **228**: 114–118.
- Bonsal BR, Zhang X, Vincent LA, Hogg WD. 2001. Characteristics of daily and extreme temperatures over Canada. *Journal of Climate* **14**: 1959–1976.
- Calderini DF, Savin R, Abeledo LG, Reynolds MP, Slafer GA. 2001. The importance of the period immediately preceding anthesis for grain weight determination in wheat. *Euphytica* **119**: 199–204.
- Carter TR. 1998. Changes in the thermal growing season in Nordic counties during the past century and prospects for the future. *Agricultural and Food Science in Finland* **7**: 161–179.
- Dhakhwa GB, Campbell CL. 1998. Potential effects of differential day-night warming in global climate change on crop production. *Climatic Change* **40**: 647–667.
- Downes P. 2007. Las heladas causaron pérdidas en cultivos de maíz, trigo y soja en el sudoeste de Buenos Aires, 24 de noviembre de 2007, Clarín Rural, <http://www.clarin.com/suplementos/rural/2007/11/24/r-01547876.htm>.
- Easterling DR. 2002. Recent changes in frost days and the frost-free season in the United States. *Bulletin of the American Meteorological Society* **83**: 1327–1332.
- Eitzinger J, Thaler S, Orlandini S, Nejedlik P, Kazandjiev V, Sivertsen TH, Mihailovic D. 2009. Applications of agroclimatic indices and process oriented crop simulation models in European agriculture. *Idojaras* **113**(1–2): 1–12.

- Feng S, Hu Q. 2004. Changes in the agro-meteorological indicators in the contiguous United States: 1951–2000. *Theoretical and Applied Climatology* **78**: 247–264.
- Fernández-Long ME, Barnatán IE, Spescha LB, Hurtado RH, Murphy GM. 2005. Caracterización de las heladas en la Región Pampeana y su variabilidad en los últimos 10 años. *Revista Facultad de Agronomía* **25**(3): 247–257.
- Fernández-Long ME, Müller GV. 2006. Annual and monthly trends in frost days in the Wet Pampa. In *Proceeding, 8th International Conference on Southern Hemisphere Meteorology and Oceanography*, 24–28 April 2006, Foz do Iguazú, Brasil.
- Forte Lay J, Scarpati OE, Capriolo A. 2008. Precipitation variability and soil water content in pampean flatlands (Argentina). *Geofísica Internacional* **47**: 341–354.
- Fraisse CW, Whidden A. 2010. Chill accumulation monitoring and forecasting. Circular EDIS AE 452: University of Florida – EDIS, <http://edis.ifas.ufl.edu/AE452>.
- Frich P, Alexander LV, Della-Marta P, Gleason B, Haylock M, Tank K., Teteron T. 2002. Observed coherent changes in climatic extremes during the second half of the twentieth century. *Climate Research* **19**: 193–212.
- Grassini P, Thorburn J, Burr C, Cassman KG. 2011. High-yield irrigated maize in the Western U.S. Corn Belt: I. On-farm yield, yield potential, and impact of agronomic practices. *Field Crops Research* **120**: 142–150, DOI: 10.1016/j.fcr.2010.09.012.
- Grau HR, Gasparri NI, Aide TM. 2005. Agriculture expansion and deforestation in seasonally dry forests of northwest Argentina. *Environmental Conservation* **32**: 140–148.
- Hall AJ, Rebella CM, Ghersa CM, Culot JPH. 1992. Field crops systems of the Pampas. In *Field Crops Systems: Ecosystems of the World*, Pearson, CJ (ed). Elsevier: Holland, 18: 413–449.
- Hall AJ, Sadras V. 2009. Whither crop physiology? In *Crop Physiology Applications for Genetic Improvement and Agronomy*, Chapter 21, Sadras VO, Calderini DF (eds). Elsevier: Amsterdam, Netherlands, 545–570.
- Haylock MR, Peterson TC, Alves LM, Ambrizzi T, Anunciacao YMT, Baez J, Barros VR, Berlato MA, Bidegain M, Coronel G, Corradi V, Garcia VJ, Grimm AM, Karloy D, Marengo JA, Marino MB, Moncunill DF, Nechoet D, Quintana J, Rebello E, Rusticucci M, Santos JL, Trebejo I, Vincent LA. 2006. Trends in total and extreme South American rainfall in 1960–2000 and links with sea surface temperature. *Journal of Climate* **19**: 1490–1512.
- Heino R, Brazdil R, Forland E, Tuomenvirta H, Alexandersson H, Beniston M, Pfister C, Rebetez M, Rosenhagen G, Rosner S, Wibig J. 1999. Progress in the study of climate extremes in northern and central Europe. *Climatic Change* **42**: 151–181.
- Hoffmann J, Nuñez S, Vargas W. 1997. Temperature, humidity and precipitation variations in Argentina and the adjacent sub-antarctic region during the present century. *Meteorological Zeitschrift, NF* **6**: 3–11.
- Hurtado RH. 2008. Relación entre el rendimiento del cultivo de trigo y el almacenaje estimado de agua en el suelo, en las cinco provincias pampeanas. Tesis de maestría en Meteorología Agrícola, Universidad de Buenos Aires, 117.
- Jones PD, Briffa KR. 1995. Growing season temperatures over the former Soviet Union. *International Journal of Climatology* **15**: 943–959, DOI: 10.1002/joc.3370150902.
- Jones PD, Briffa KR, Osborn TJ, Moberg A, Bergström H. 2002. Relationships between circulation strength and the variability of growing season and cold season climate in northern and central Europe. *Holocene* **12**: 643–656.
- Katz RW, Brown BG. 1992. Extreme events in a changing climate: variability is more important than averages. *Climate Change* **21**: 289–302.
- Kendall MG. 1975. *Rank Correlation Methods*, Oxford University Press: New York.
- León AJ, Lee M, Andrade FH. 2001. Quantitative trait loci for growing degree days to flowering and photoperiod response in Sunflower (*Helianthus annuus* L.). *Theoretical and Applied Genetics* **102**: 497–503, DOI: 10.1007/s001220051673.
- Liebmann B, Dole RM, Jones C, Bladé I, Allured D. 2010. Influence of choice of time period on global surface temperature trend estimates. *Bulletin of the American Meteorological Society* **91**: 1485–1491, DOI: 10.1175/2010BAMS3030.1.
- Lobell DB. 2007. Changes in diurnal temperature range and national cereal yields. *Agric Forest Meteorol* **145**: 229–238, DOI: 10.1016/j.agrformet.2007.05.002.
- Magra G, Senn J, Martín R. 2005. Evaluación del daño provocado por heladas tardías en trigo. *Revista Agromensaje* **17**: 2005, <http://www.fcagr.unr.edu.ar/Extension/Agromensajes/17/15AM17.html>.
- Magrin GO, Díaz RA, Travasso MI, Rodriguez G, Boullón D, Nuñez M, Solman S. 1998. Vulnerabilidad y mitigación relacionada con el impacto del cambio global sobre la producción agrícola. In *Proyecto de Estudio sobre el Cambio Climático en Argentina* Barros, V, Hoffmann, JA, Vargas, WM (eds). Proyecto ARG/95/G/31-PNUD-SECYT, Buenos Aires, Argentina, 290.
- Magrin GO, Travasso MI. 2002. An integrated climate change assessment from Argentina. In *Effects of Climate Change and Variability on Agricultural Production Systems*, Chapter 10, Randolph, JC, Doering, O, Southworth, J, Pfeifer, R (eds). Kluwer Academic Press: Boston, 193–218.
- Magrin GO, Travasso MI, Rodriguez GR. 2005. Changes in climate and crop production during the 20th century in Argentina. *Climatic Change* **72**: 229–249, DOI: 10.1007/s10584-005-5374-9.
- Mall RK, Gupta BRD, Singh KK, Singh TK. 2000. Phenology and yield of wheat (*Triticum aestivum*) in relation to growing degree-days and photothermal units. *Indian Journal of Agricultural Sciences* **70**: 647–652.
- Mann HB. 1945. Non-parametric tests against trend. *Econometrica* **13**: 245–259.
- Menzel A, Jakobi G, Ahas R, Scheifinger H, Estrella N. 2003. Variations of the climatological growing season (1951–2000) in Germany compared with other countries. *International Journal of Climatology* **23**: 793–812, DOI: 10.1002/joc.915.
- Minetti JL, Vargas WM. 1983. Comportamiento del borde anticiclónico subtropical en Sudamérica. I Parte. *Meteorológica* **14**: 645–655.
- Minetti JL, Vargas WM. 1992. Comportamiento del borde anticiclónico subtropical en Sudamérica. II Parte. *Revista Geofísica (IPGH.OEA)* **33**: 179–190.
- Minetti JL, Vargas WM, Poblete AG, Acuna LR, Casagrande G. 2003. Non-linear trends and low frequency oscillations in annual precipitation over Argentina and Chile, 1931–1999. *Atmosfera* **16**: 119–135.
- Monzón JP, Sadras VO, Abbate PA, Caviglia OP. 2007. Modelling management strategies for wheat–soybean double crops in the south-eastern Pampas. *Field Crops Research* **101**: 44–52, DOI: 10.1016/j.fcr.2006.09.007.
- Moonen AC, Ercoli L, Mariotti M, Masoni A. 2002. Climate change in Italy indicated by agrometeorological indices over 122 years. *Agricultural and Forest Meteorology* **111**: 13–27, DOI: 10.1016/S0168-1923(02)00012-6.
- Myneni RB, Keeling CD, Tucker CJ, Asrar G, Nemani RR. 1997. Increased plant growth in the northern high latitudes from 1981 to 1991. *Nature* **386**: 698–702, DOI: 10.1038/386698a0.
- Nakamura H, Lin G, Yamagata T. 1997. Decadal climate variability in the North Pacific during the recent decades. *Bulletin of the American Meteorological Society* **78**: 2215–2225.
- Nuñez MN, Ciapessoni HH, Rolla A, Kalnay E, Cai M. 2008. Impact of land use and precipitation changes on surface temperature trends in Argentina. *Journal of Geophysical Research* **113**: 1–11, DOI: 10.1029/2007JD008638.
- Otegui ME, Ruiz RA, Petrucci D. 1996. Modelling hybrid and sowing date effects on potential grain yield of maize in a humid temperate region. *Field Crops Research* **47**: 167–174, DOI: 10.1016/0378-4290(96)00031-7.
- Otegui ME, Lopez Pereira M. 2003. Fecha de siembra. In *Producción de Granos. Bases Funcionales Para su Manejo*, Satorre E, Benech-Arnold RL, Slafer GA, de la Fuente E, Miralles D, Otegui ME, Savín R (eds). Editorial Facultad de Agronomía – Universidad de Buenos Aires: Buenos Aires, 259–275.
- Pascale AJ, Damario EA. 2004. *Bioclimatología Agrícola y Agroclimatología*. Editorial Facultad de Agronomía – Universidad de Buenos Aires: Buenos Aires, 550.
- Peterson TC, Folland CK. 2000. Climate change detection what can a small working group do about it? *CLIVAR Exchanges* **5**: 24–25.
- Quiroz F, Maneiro C. 2007. Informe sobre heladas y daño en girasol. Informe Técnico EEA INTA Balcarce, diciembre 2007, <http://www.inta.gov.ar/balcarce/noticias/2007/heladasgirasol.htm>.
- Raper CD, Kramer PJ. 1987. Stress physiology. In *Soybeans: Improvement, Production and Uses* Wilcox JR (ed). Am. Soc. of Agron.: Madison, WI.
- Rinaldi M. 2004. Water availability at sowing and nitrogen management of durum wheat: a seasonal analysis with the CERES-Wheat model. *Field Crops Research* **89**: 27–37, DOI: 10.1016/j.fcr.2004.01.024.
- Root TL, Price JT, Hall KR, Schneider SH, Rosenzweig C, Pounds JA. 2003. Fingerprints of global warming on wild animals and plants. *Nature* **421**: 57–60, DOI: 10.1038/nature01333.

- Ross F, Carpaneto B, Andrade F, Quiroz F, Maneiro C. 2007. Boletín informativo sobre daño por heladas. INTA, <http://www.inta.gov.ar/actual/info/boletinheladas.pdf>.
- Rosenzweig C, Tubiello FN. 1996. Effects of changes in minimum and maximum temperature on wheat yields in the central US: a simulation study. *Agricultural and Forest Meteorology* **80**: 215–230, DOI: 10.1016/0168-1923(95)02299-6.
- Rusticucci M, Penalba O. 2000. Interdecadal changes in the precipitation seasonal cycle over Southern South America and their relationship with surface temperature. *Climate Research* **16**: 1–15, DOI: 10.3354/cr016001.
- Rusticucci M, Barrucand M. 2004. Observed trends and changes in temperature extremes over Argentina. *Journal of Climate* **17**: 4099–4107.
- Salinger MJ, Griffiths GM. 2001. Trends in New Zealand daily temperature and rainfall extremes. *International Journal of Climatology* **21**: 1437–1452, DOI: 10.1002/joc.694.
- Schwartz MD, Rein A, Anto A. 2006. Onset of spring starting earlier across the northern hemisphere. *Global Change Biology* **12**: 343–351, DOI: 10.1111/j.1365-2486.2005.01097.
- Sen PK. 1968. Estimates of the regression coefficient based on Kendall's Tau. *Journal of the American Statistical Association* **63**: 1379–1389.
- Shen SSP, Yin H, Cannon K, Howard A, Chetner S, Karl TR. 2005. Temporal and spatial changes of the agroclimate in Alberta, Canada, from 1901 to 2002. *Journal of Applied Meteorology* **44**: 1090–1105, DOI: 10.1175/JAM2251.1.
- SIIA 2010. *Sistema Integrado de Información Agropecuaria. Ministerio de Agricultura, Ganadería y Pesca*. Presidencia de la Nación: Argentina, <http://www.siiia.gov.ar/index.php>.
- Slafer GA, Rawson HM. 1994. Sensitivity of wheat phasic development to major environmental factors: A re-examination of some assumptions made by physiologists and modellers. *Australian Journal of Plant Physiology* **21**: 393–426, DOI: 10.1071/PP9940393.
- Song Y, Linderholm HW, Chenb D, Walther A. 2010. Trends of the thermal growing season in China, 1951–2007. *International Journal of Climatology* **30**: 33–43, DOI: 10.1002/joc.1868.
- Spescha L. 2008. *Relación entre la reserva del agua en el suelo y los rendimientos de soja de la región oriental de secano de la Argentina*. Tesis de Maestría de Meteorología Agrícola: Universidad de Buenos Aires: Buenos Aires, 120.
- Trenberth KE, Hurrell JW. 1994. Decadal atmosphere-ocean variations in the Pacific. *Climate Dynamics* **9**: 303–319, DOI: 10.1007/BF00204745.
- Trenberth KE, Jones PD, Ambenje P, Bojariu R, Easterling D, Klein Tank A, Parker D, Rahimzadeh F, Renwick JA, Rusticucci M, Soden B, Zhai P. 2007. Observations: surface and atmospheric climate change. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* Solomon, S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds). Cambridge University Press: Cambridge and New York, NY.
- Tubiello FN, Rosenzweig C, Goldberg RA, Jagtap S, Jones JW. 2002. Effects of climate change on US crop production: simulation results using two different GCM scenarios. Part I: wheat, potato, maize, and citrus. *Climate Research* **20**: 259–270, DOI: 10.3354/cr020259.
- Tucker CJ, Slayback DA, Pinzon JE, Los SO, Myneni RB, Taylor MG. 2001. Higher northern latitude normalized difference vegetation index and growing season trends from 1982–1999. *International Journal of Biometeorology* **45**: 184–190, DOI: 10.1007/s00484-001-0109-8.
- Vincent LA, Peterson T C, Barros, VR, Marino MB, Rusticucci M, Carrasco G, Ramirez E, Alves LM, Ambrizzi T, Berlatto MA, Grimm AM, Marengo JA, Molion L, Moncunill DF, Rebello IE, Anunciação YMT, Quintana J, Santos JL, Baez J, Coronel G, Garcia J, Trebejo I, Bidegain M, Haylock MR, Karoly D. 2005. Observed trends in indices of daily temperature extremes in South America 1960–2000. *Journal of Climate* **18**: 5011–5023, DOI: 10.1175/JCLI3589.1.
- Walther A, Linderholm HW. 2006. A comparison of growing season indices for the Greater Baltic Area. *International Journal of Biometeorology* **51**: 107–118, DOI: 10.1007/s00484-006-0048-5.
- Walther GR, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin JM, Hoegh-Guldberg O, Bairlein F. 2002. Ecological responses to recent climate change. *Nature* **416**: 389–395, DOI: 10.1038/416389a.
- Wang L, Chen W, Huang RH. 2007. Changes in the variability of North Pacific Oscillation around 1975/1976 and its relationship within East Asian winter climate. *Journal of Geophysical Research* **112**: D11110, DOI: 10.1029/2006JD008054.
- Wang L, Chen W, Zhou W, Chan JCL, Barriopedro D, Huang R. 2010. Effect of the climate shift around mid 1970s on the relationship between wintertime Ural blocking circulation and East Asian climate. *Journal of Climatology* **30**: 153–158, DOI: 10.1002/joc.1876.
- Yaodong DU. 2005. Frost and high temperature injury in China. In *Natural Disasters and Extreme Events in Agriculture* Sivakumar MVK, Motha RP, Das HP (eds). Springer: Netherlands, 145–157.
- Zhang X, Zwiers FW. 2004. Comments on “Applicability of prewhitening to eliminate the influence of serial correlation on the Mann-Kendall test. *Water Resources Research* **40**: W03805, DOI: 10.1029/2003WR002073.