

Innovations for a sustainable future: rising to the challenge of nitrogen greenhouse gas management in Latin America

Mercedes MC Bustamante¹, Luiz A Martinelli²,
Jean PHB Ometto³, Janaina Braga do Carmo⁴,
V́ctor Jaramillo⁵, Mayra E Gavito⁵, Patricia I Araujo⁶,
Amy T Austin⁶, Tibisay Ṕrez⁷ and Sorena Marquina⁷



Latin America encompasses a dizzying array of ecosystems and socioeconomic models, and the region will be highly vulnerable to the projected impacts of climate change in the next century. At the same time, Latin America can significantly contribute to the mitigation of greenhouse gases (GHG) emissions within a sustainable development framework. Land use conversion with associated biomass burning, agriculture with N fertilizers and animal waste are the main anthropogenic sources of nitrous oxide (N₂O) emissions in the region, and have increased markedly in the last decades. Effective sustainable management for the mitigation of N₂O emissions requires the proper evaluation of all sources, many of which are still roughly estimated or unknown, testing alternatives to reduce primary sources, and technological innovation for higher resource-use efficiency within the farm. Current barriers might be overcome through policies that support sustainable practices that reduce negative environmental impacts and simultaneously maintaining ecosystem function and services.

Addresses

¹ Departamento de Ecologia, Universidade de Braślia, Braślia CEP 70910-900, Brazil

² CENA-Universidade de S~o Paulo, Avenida Centen~rio 303, 13416-000 Piracicaba, SP, Brazil

³ Centro de Ci~ncia do Sistema Terrestre. Instituto Nacional de Pesquisas Espaciais, Av. dos Astronautas, 1758 S~o Jos~ dos Campos, SP, Brazil

⁴ Universidade Federal de S~o Carlos, Campus Sorocaba, Rodovia Jos~ Leme dos Santos, km 110, SP 264, Itinga, 18052-780 Sorocaba, SP, Brazil

⁵ Centro de Investigaciones en Ecosistemas, Universidad Nacional Aut3noma de M~xico A.P. 27-3 Santa Mar~a de Guido, C.P. 58090, Morelia, Michoac~n, M~xico.

⁶ IFEVA-CONICET, Universidad de Buenos Aires, Buenos Aires C1417DSE, Argentina

⁷ Instituto Venezolano de Investigaciones Científicas, Caracas, Venezuela

Corresponding author: Bustamante, Mercedes MC (mercedes@unb.br, mercedesmcb@gmail.com)

Current Opinion in Environmental Sustainability 2014, 9–10:73–81

This review comes from a themed issue on **System dynamics and sustainability**

Edited by **Carolien Kroeze**, **Wim de Vries** and **Sybil Seitzinger**

For a complete overview see the [Issue](#) and the [Editorial](#)

Received: 10 March 2014; Accepted: 9 September 2014

Available online 29th September 2014

<http://dx.doi.org/10.1016/j.cosust.2014.09.002>

1877-3435/© 2014 Elsevier B.V. All rights reserved.

Introduction

A recent analysis of the challenges for Latin America and the Caribbean (LAC) region related to physical and socioeconomic impacts of climate change concluded that the region is particularly vulnerable to the observed and projected effects of climate change due to its geographic location, population distribution, infrastructure, and reliance on fragile or non-renewable natural resources for economic activities and livelihood [1*]. The conservative projection of yearly economic damages in LAC caused by some of the major physical impacts associated with the projected rise of 2°C in global mean air temperature is approximately 2.2% of the region's 2010 gross domestic product (GDP, \$4.6 trillion). Potential losses of this magnitude clearly undermine the region's prospects for improvements in the quality of life by significantly limiting development options and severely restricting access to natural resources and ecosystem services, all with socially damaging consequences for equity and poverty levels [1*].

While the region shows genuine vulnerability to the projected impacts of climate change, LAC can also significantly contribute to the mitigation of greenhouse gases (GHGs) emissions within a sustainable development framework. Latin America accounted for 8% of the world's GHG emissions in 2005 [2], which include nitrous oxide (N₂O), carbon dioxide (CO₂) and methane (CH₄). Although N₂O is naturally present in the atmosphere as part of the Earth's nitrogen cycle, human activities are increasing the amount of atmospheric N₂O, particularly through agricultural activities. Specifically in Latin America, land use change (biomass burning — 50%), agriculture (N-fertilizers — 10%) and animal waste (40%) are the main sources of N₂O anthropogenic emissions [3]. Such increase is of particular concern due to the radiative forcing potential of N₂O (300 times that of CO₂ over a 100-year timescale on a per mole basis) and the strong correlation with increased emissions and agricultural intensification in the region.

Recently, the impacts of changes on the regional N cycle in Latin America were evaluated [4*]. These authors highlighted the lack of detailed information on many aspects of the nitrogen cycle, which is a serious impediment to our ability to evaluate and project how human


activity is altering nitrogen pools and turnover at regional scales. Here, we expand this assessment to focus on the main drivers involved in human activities associated with increased GHGs and specifically nitrous oxide (N₂O) emissions in Latin America. In addition, we highlight potential mitigation strategies and projections for research needs and priorities.

Regional GHG emissions by sectors

Brazil contributed half of all regional GHG anthropogenic emissions in 2005 and together with Mexico, Venezuela and Argentina, accounted for nearly 80% of total emissions [2]. The share of total emissions among economic sectors is more critical when considering potential mitigation measures that could be adopted in the region. In 2005, the sectors 'Land use changes and forestry (LUCF)' and 'Agriculture' contributed with largest shares of total anthropogenic GHG emissions across the region, with 47% and 20% respectively (Figure 1). The Venezuelan national inventory, however, showed differences in the relative emissions by sector. The energy sector represented the largest source of GHG (75% of total emissions) followed by agriculture (17%) while LUCF represented a *net sink* of approximately 14 300 Gg CO₂ eq [5] (MARN-Venezuela, 2005). In addition N₂O emissions country-level emissions ranged widely, with 3% (Mexico) to 28% (Argentina) of total anthropogenic GHG emissions for the region (www.unfccc.int/ghg_data_unfccc).

GHG emissions from LUCF also showed important changes in other countries. Brazil's share of regional and global GHG emissions from land use changes was particularly significant in 2005, but has declined in the last decade. A recent update of emissions figures [6] indicate that in 2010, Brazil had reduced GHG emissions by nearly half, to 1.25 Pg CO₂ eq, compared to baseline emissions of 2.03 Pg CO₂ eq in 2005, strongly associated with the reduction of deforestation rates in the Amazon basin. In Argentina, between 1990 and 2000 [7], the LUCF sector showed greater relative changes, however, but with an increase in the net carbon sink of nearly 200%. Native

Figure 1



Contributions of total anthropogenic greenhouse gas emissions in Latin America and the Caribbean (LAC) from different sources, 2005. The contributions refer to percentage shares of total anthropogenic GHG emissions from LAC, and not the total fraction of each sector's (i.e. energy) contribution.

Source: Vergara [1*] based on data compiled from World Resources Institute (2012).

vegetation conversion to agriculture and ranching also decreased in Mexico between 1990 and 2010 [8], which probably contributed to decreased GHG emissions in this country as well.

The comparison of N₂O emissions by country (Table 1) indicates that Brazil is by far the largest emitter in the region. In general, agriculture accounts for the largest share of N₂O emissions in the region (up to 96.8%). Globally, N₂O emission from agriculture is equivalent to about 66% of total gross anthropogenic emissions [9]. Agriculture plays a key role in the LAC economies, accounting for approximately 6% of regional gross domestic product (GDP) and 15% of employment in


Table 1

Total anthropogenic emissions of greenhouse gases (GHG, Tg CO₂ eq) and N₂O (Tg CO₂ eq) from the largest emitters in Latin America. Data based on National Inventories of GHG submitted to the UNFCCC and prepared according IPCC guidelines

Country	Argentina		Brazil		Mexico		Venezuela
	2000	2000	2005	2000	2006	1999	
Total GHG	238.70	2087.66	2191.86	563.23	711.65	177.90	
Total N ₂ O (% of total GHG)	67.56 (28.3%)	169.20 (8.1%)	546.00 (24.9%)	12.13 (2.1%)	20.51 (2.9%)	16.15 (9.1%)	
N₂O by sectors (% of total N₂O)							
Agriculture	65.39 (96.8%)	121.68 (71.9%)	476.20 (87.2%)	7.46 (61.5%)	6.99 (34.1%)	15.42 (95.5%)	
Land use change	0.06	6.45	20.90	0.31	0.17	0.01	
Energy	1.01	2.98	12.10	2.50	10.95	0.22	
Industry	0.15	6.17	22.80	0.11	0.36	0.08	
Waste	0.96	3.84	14.00	1.96	2.05	0.42	

Source: www.unfccc.int/ghg_data_unfccc, July 18, 2014.

Figure 2



A general overview of different sources of N₂O-N emissions (Gg yr⁻¹) in the LAC region. Bars indicate total emissions per year calculated for the years of reference. The different colors for agricultural soils indicate the range of estimates for emissions from this sector. Data for agricultural soils (croplands with N fertilizer and application of manure) from [25,26]; biomass burning associated to deforestation [11]; animal waste [11] and biomass burning for fuels [16,18,19].

2010 [4^{*}]. Agriculture also represents a key factor in regional food security. In the beginning of the 1960s, arable land in LAC was responsible for 7% of the global arable land area; 50 years later this proportion has increased to 11% [10^{*}]. More than 50% of the global cultivation of sugarcane and coffee occur in Latin America, while soybean occupies more than 40% of the total global area [10^{*}]. The use of N-fertilizer has grown from only 5 kg ha⁻¹ to approximately 50 kg ha⁻¹ over the last fifty years [10^{*}].

Biomass burning (from landscape fires, household cooking and heating) represents about 11% of total gross anthropogenic global N₂O emissions [9]. In particular, Latin America alone emitted nearly 40% of the global N₂O emissions due to biomass burning in 1990, a disproportional contribution considering that Latin America corresponds to only about 13% of global land surface [11].

The following sections provide a more detailed perspective on the regional emissions of N₂O related to biomass burning and agriculture with a synthesis of available data from different sources (Figure 2). Despite the differences in source years and data sources, we attempted to assess the overall global relevance of these activities in the region and the potential to mitigate N₂O emissions. Clearly, the relative importance of agricultural activity is highlighted here, but additionally, the underappreciated role of biomass burning for contributing to N₂O emissions is approaching agriculture in terms of its relative contribution to overall emissions.

N₂O emissions from land use change and biomass burning

The clearing of natural vegetation, burning, fertilization of agricultural lands, intensive cattle ranching and increasing dominance by legume species in areas under secondary succession, have been identified as causes of increasing N₂O and NO emissions in tropical regions (see [12] for a review). However, large uncertainties remain for regional estimates of trace gas fluxes in Latin American ecosystems due to the scarcity of data with adequate spatial distribution and a combination of social and ecological factors that may affect the fluxes at local scale [4^{*}].

Deforestation and biomass burning are important sources of N₂O emissions, but vary substantially among the major biomes in LAC. More than 80% of deforestation occurs in humid and dry forest, and savannas/shrublands [13^{*}], and these conversions are most often accompanied by burning of extant vegetation or debris. It is estimated that from 15 to 30 million ha are burned per year in LAC [14,15]. The N₂O emissions caused by biomass burning associated with deforestation in 1990 were estimated to be near 200 Gg of N-N₂O in the region [11]. Subtropical regions in South America are also indirectly affected by biomass burning in tropical areas due to the long-range transport of smoke from Amazonia to the southern and southeastern part of the South American continent [17]. Fuel use for cooking and home heating is an additional source of burning, and it is estimated that 350 million tons of wood are burned per year in LAC, with half this amount in Brazil alone [16]. Assuming that 6.4 g N-N₂O is produced per Giga Joule (GJ) of fuel-wood burned [18] and that 1 ton of fire-wood produces 13 GJ of energy [19], we estimate that 350 million tons of wood burned would emit approximately 30 Gg of N-N₂O, which is currently not incorporated into any estimate of emissions from the region.

Following deforestation, most areas are converted to agriculture and grazing (pasture) lands [20]. The majority of natural vegetation areas that are burned are related to the opening of new grazing and agriculture areas. For example, Argentina experienced unprecedented deforestation in dry forests between 1977 and 2008 as a consequence of agricultural expansion [21], particularly for soybean production [22,23]. Paruelo *et al.* [21] estimated that the burning of 8.7 million tons of biomass yr⁻¹ due to deforestation fires in Argentina produced emissions of 0.2 Gg of N₂O. Between 1994 and 2000, GHG emissions, mostly due to the burning of sugar cane residues, ranged between 187 and 208 Gg CO₂ eq (20% was from N₂O) [7]. Biomass burning has also increased in Mexico during the past 40 years although there is substantial inter-annual variability [24]; burning of agricultural residues is common in rural landscapes. Current numbers, based mainly on sugar cane cropping, account for only 9 Gg CO₂ eq [7] emissions per year,

which is likely to be a serious underestimate of this source. In Venezuela [5], burning of agricultural residues and savanna areas emitted 0.1% and 0.4% of total N₂O emissions. On the basis of satellite measurements, it is likely that emissions from savanna burning are underestimated considering the dramatic increase in burned area in recent years in LAC [14]. As well, these authors have suggested an enhanced relative importance of small fires from savannas and grasslands to the global GHG emissions, all of which are currently undocumented.

N₂O emissions from agriculture

N₂O emissions from agriculture include direct emissions from agricultural soils, principally due to the application of animal manure and mineral N-fertilizers, and manure production in pastures. They also include indirect emissions resulting from the subsequent leaching of nitrate to ground water and surface waters, and from ammonia deposition that had volatilized as a result of agricultural activities. Estimates of ammonia (NH₃) volatilization and nitrate (NO₃⁻) leaching for Latin America in comparison to world values for the period between 1970 and 2030 are presented in Table 2. There is a consistent increase in both NH₃ volatilization and nitrate (NO₃⁻) in the region in absolute and relative values, which is associated with increased agricultural intensity in both the crop and livestock sector. These trends suggest that indirect emissions of N₂O also will increase in the future.

For South and Central America, it was estimated that agricultural soils in 2005 emitted approximately 240 Gg of

N-N₂O [25] up to approximately 360 Gg of N-N₂O [26] (Figure 2) due to the use of mineral N-fertilizers, to crop residues and animal manure [25]. This is equivalent to almost 10% of the global emissions, while arable land in LA is equivalent to 15% of the global estimate.

The use of nitrogen fertilizers and other agronomic inputs in agriculture is imbalanced in the Latin American region, due to socio-economic factors that limit access to these inputs for many small landholders, combined with ecological factors, including baseline natural fertility of agricultural systems [4]. For example, Argentina consumes 60% of the fertilizer in the Southern Cone countries (Argentina, Chile, Paraguay, and Uruguay) while in Brazil, three crops (maize, soybean and sugar cane) are responsible for 56% of the N, 71% of the P₂O₅ and 75% of the K₂O fertilizer application. Most of the emissions of GHGs in the agricultural sector in Brazil are associated with the domestic livestock and the cultivation of soybean, maize, sugar cane and rice, which together occupy more than 70% of the country's cultivated area. Emissions in this agricultural sector increased 37% from 1990 to 2005, considering primarily CH₄ and N₂O. In this period, crop productivity increased well above the extension of land area used for agricultural production [27]. In contrast to Brazil, CH₄ and N₂O emissions in Mexico have remained stable during the period 1990–2010, with CH₄ representing 43% and N₂O 57% of emissions [28]. In 2010, N₂O emissions from soil management (~150 Gg N₂O) represented the largest contributor to GHG emissions from the agricultural sector in Mexico, likely attributed to the use of mineral N-fertilizers resulting from production. In Argentina, GHG emissions in the agricultural and cattle sector during the year 2000 represented 43% of total GHGs; 21% was associated with agricultural practices (primarily N₂O emissions) and 22% with livestock production (mostly CH₄) [29]. Direct N₂O emissions from agricultural lands increased 85% between 1990 and 2000 (from ~63.2 to 117.2 Gg N yr⁻¹), in line with the increment of nitrogen incorporated to the soil (from ~3.3 to 5.6 Gg N yr⁻¹), mainly due to cultivation of leguminous species, but also from mineral N-fertilizers [7]. While emission from synthetic nitrogen fertilization was 2% of the total N₂O emission in 1990, this value climbed to 12% by 2000 [29]. In Venezuela, total N₂O emissions from agriculture (48 Gg N₂O) have three major sources: direct emissions from agricultural soils (15 Gg N₂O); soils in grazed pastures (19 Gg N₂O); and indirect emissions from leaching and runoff (12 Gg N₂O) [5]. Recently measured N₂O emission factors (EF) derived from Venezuelan agricultural soils show a large range (0.30–6.1% of the applied N fertilizer), with overall average values of 1.9% higher than IPCC default value (1%) [30].

Increased biofuel production has been associated with direct and indirect land-use change, changes in land

Table 2

Comparison of total ammonia (NH₃) volatilization (includes NH₃-N emissions from fertilizer and animal manure application, grazing, and from animal housing and manure storage systems). N₂O and NO emissions are based on fertilizer and animal manure application (excluding emissions from fallow land), and nitrate leaching for intensive agricultural systems (NO₃-N, Tg yr⁻¹)^a for Latin America. Percent values indicate the share of Latin America contribution to the world values

Region	Total NH ₃ -N	N ₂ O-N	NO-N	NO ₃ -N
1970				
Latin America	1.6	0.2	0.1	1.0
World	18.1	2.0	1.1	18.2
(%)	(8.8)	(10.0)	(9.1)	(5.5)
1995				
Latin America	3.7	0.3	0.1	2.9
World	34.2	2.7	1.5	28.5
(%)	(10.8)	(11.1)	(6.7)	(10.2)
2030				
Latin America	5.6	0.5	0.2	4.4
World	44.0	3.5	2.0	35.3
(%)	(12.7)	(14.3)	(10.0)	(12.5)

Data from Bowman *et al.* [49].

^a Intensive agricultural systems include total arable land and grassland in mixed/industrial livestock production systems; pastoral systems are excluded.

management practices, and increased application of fertilizers and pesticides. In Latin America, bioenergy development is based on two major crops: sugarcane and soybean. In 2008, Cruzten *et al.* [31] established that if the emission factor of N₂O in sugarcane crops surpasses 5% of the quantity of nitrogen in the fertilizers, the environmental damages caused by N₂O emissions would not offset the carbon gain by the biofuel displacement of fossil fuels. Recent studies based on field measurements in Brazil indicated that the emission factor associated with N-fertilizer use was well below 3%. Practices which include the joint application of vinasse and fertilizer, and associated with a large amount of crop straw remaining in the soil, can result in an emission factor of nearly 3% [32]. Sugarcane crop residues from green cane ('non-burning') harvests also increase N₂O emission when soil moisture increases [33].

Soybean cultivation is a very important activity in the agricultural sector of both Argentina and Brazil [34] and the rapid increase in areas devoted to this crop will have important consequences for N cycling in the region. Although emissions from soybean fields are considered low in Brazil [35] and Argentina [36] relative to N-fertilized crops, the large extension of cultivated area (about 25 million hectares in Brazil and 19 million hectares in Argentina, in 2012) with this crop might represent a significant source of N₂O emissions simply due to the very large area currently under cultivation. Very little is known regarding the potential source of N₂O emissions from this important agricultural activity in the region and warrants further research.

Livestock and manure management

The importance of Latin America in the livestock sector is also considerable, since the region hosts approximately 20% of the global cattle population and a similar proportion of poultry [9]. Although this is an underappreciated source of N₂O emissions, estimates for the region's N-N₂O emissions in 1990 from livestock waste were nearly 180 Gg of nitrogen [10*]. This value is equivalent to almost 20% of the global livestock related N-N₂O emissions for the same year. While later estimates did not distinguish emissions derived from mineral N-fertilizers from those derived from animal manure [25,26], it was estimated that approximately 3400 Gg of N as manure were used in LA [26], which is comparable to the 5700 Gg of N applied to croplands as N-fertilizer.

This is particularly relevant for the region as N₂O emissions related to cattle excreta in grazed pastures is considered 33% higher in South America than for the globe on average (23 kg CO₂ eq kg⁻¹ vs. 17 kg CO₂ eq kg⁻¹, respectively). This is due to the fact that cattle production in LA is largely pasture-based with open ranges for the animals, and as such, the animals increase in mass more slowly and manure deposited in pasture is more prone to

N₂O formation than in feed lots [37]. Nevertheless, a recent field study in central Brazil reported a direct emission factor (EF) for N₂O of 0.007 for the cattle excreta as whole, well below the IPCC EF (0.02) [38].

N₂O emissions from livestock and manure management in Mexico have also remained relatively stable at about 6445 Gg CO₂ eq between 1990 and 2010 due to an overall stable livestock population: cattle rearing has actually decreased, but with a simultaneous increase in poultry production [28]. In Argentina, direct emissions of N₂O from cattle excreta are the second source of GHG emission of livestock sector, being 18 300 Gg CO₂ eq in the year 2000 (21% of the total GHG emission of the sector). Indirect emissions from N volatilization from cattle excreta and urine were 8940 Gg CO₂ eq (10.4% of the total Gg emissions) and relatively stable during the period 1990–2000. Emissions from manure management played a minor role, accounting for 0.2% of the total emissions of the sector in the year 2000 [7].

Mitigation options – innovation and sustainable future

The Global Forest Resources Assessment [39] estimated that Latin America has suffered the largest net loss of forests in the world from 1990 to 2010. South America alone lost approximately 80 million ha of forests in these two decades, which corresponds to a deforestation rate of 4 million ha per year [39]. Forest fragmentation and the use of fire as a land management practice are major drivers of biomass burning and associated N₂O emissions in Latin America. Fire prevention and management (in fire-prone ecosystems such as savannas) is thus a key mitigation action. The need for proactive fire management has been also mentioned in other studies, as a substantial loss of ecosystem services [4*,27*]. Our limited understanding of the science of fire dynamics in the region hampers our assessment of the impacts of landscape fires in the face of global warming and increasing anthropogenic disturbance. In addition, the complexity of deterring deforestation and biomass burning due to the interplay of socio-economic, cultural and political drivers and the undocumented nature of many aspects of these dynamics including the importance of small fires and wood burning in homes makes it a particularly daunting challenge for mitigation options.


Given the importance of fire and unregulated biomass burning in affecting N₂O emissions in LA, it would be important to combine the monitoring of land cover with early-warning systems for identification and landscape fire prevention. One of the most efficient approaches is through the use of fire forecasting models. These models aid in identifying the main temporal, spatial and climatic factors that contribute to fire outbreaks and can therefore be employed to minimize impacts. Models to assess smoke-spread can also contribute to prior identification

of the areas potentially damaged by fire, thus supporting the decision-making process and possibly reducing the impact of such events.

Another important source of N_2O is the emission from agricultural soils under cultivation, mainly due to the use of synthetic N-fertilizers and organic sources of N in manure. As N_2O is formed principally from N that is not directly utilized by the crop, the basic principle would be to promote agricultural practices that enhance the N-use efficiency by crops, thereby avoiding N losses to the environment [40]. For instance, studies in the region indicated that N_2O emissions vary depending on the nitrogen source and application of N fertilizer [41] and chemical composition of the fertilizer [30]. This reinforces the importance of the definition of specific critical levels of N fertilizer to enhance productivity while at the same time minimizing losses due to N_2O emissions.

More efficient land management and major biological innovations in agriculture have the potential to increase productivity while decreasing environmental impacts. For example, Biological Nitrification Inhibition (BNI) is a process by which certain plants — in particular the tropical pasture grass *Brachiaria humidicola* — naturally inhibit the conversion of reactive N in the soil to forms subject to leaching (NO_3) or gaseous (N_2O) losses. These natural reductions of N losses from the soil under managed pastures have a direct and beneficial environmental effect [42]. Additionally, it has been suggested that the adoption of no-till (NT) agriculture could decrease N_2O emissions when compared with conventional tillage (CT) agriculture [43]. Recently, Van Kessel *et al.* [44] evaluated the effect of tillage on N_2O emissions in a meta-analysis comparing (CT) and NT and reduced tillage (RT) agriculture in humid and dry climatic zones. They concluded that in humid climates, deep placement of fertilizer-N is recommended in conjunction with NT or RT agriculture

Figure 3



Main drivers of N_2O emissions (agriculture and land use changes associated with biomass burning) in Latin America and potential mitigation strategies for these emissions considering feedbacks on climate change and sustainable agriculture.

to minimize N₂O emissions. Additionally, in dry climates, NT/RT practices alone are an effective mitigation strategy for reducing N₂O emissions if sustained for a prolonged time. NT practices have been adopted in several countries of Latin America [10^{*}], and in Brazil alone, there are more than 25 million ha under NT agriculture [45] with the first areas being implemented near the 1970s. In Argentina, while GHGs emissions increased over the last 50 years in those areas where high deforestation and burning occurred, emissions in the Pampas were substantially reduced as a consequence of NT practices [46].

Managing an integrated data synthesis and modeling research network for reducing N₂O emissions from agricultural soils in Latin America is crucial for effective agricultural management practices in the region. A major target is to start from a comprehensive evaluation of all GHG sources (many still roughly estimated or unknown, like compost, organic fertilizers and pesticides), testing of alternatives directed to reduce the identified main GHG sources, and technological innovation for energy production and recycling of both energy and materials at the individual farm scale. Small stakeholder agricultural practices in many regions of LAC are very important for local food production, and the impact of these practices on GHG emissions is almost entirely undocumented. In addition, current practices claiming to be environmentally friendly, including organic production systems, may be quite inefficient in energy use and produce similar or greater GHG emissions as conventional systems [47], in spite of some evidence from a global meta-analysis that suggests that larger soil carbon storage, N₂O emission reductions and larger methane uptake can be achieved [48].

Clearly, increased resource use efficiency is a major goal in Latin American countries that encompasses solutions to the problems highlighted in this review. Some relevant measures to mitigate N₂O emissions in LAC are summarized in Figure 3. Because of the clearing of land for agricultural conversion, the inevitable increase in GHG emissions generate substantial ecological imbalances that need to be restored, but this cannot occur in isolation with a single focus on reduction of GHG emissions. Current barriers might be overcome through policies that support truly sustainable practices aimed at reducing multiple negative environmental impacts, including reducing GHG emissions, and increasing the maintenance of ecosystem functions and services [4^{*}].

Acknowledgements

This study was supported by IAI-CRN3 project 'Nitrogen Cycling In Latin America: Drivers, Impacts and Vulnerabilities (Nnet)'. Additional support came from UBACyT and ANPCyT of Argentina, and the New Phytologist Trust. MMC Bustamante received financial support from CNPq (no. 312035/2013-6).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Vergara W, Rios AR, Galindo LM, Gutman P, Isbell P, Suding PH, Samaniego J: *The Climate and Development Challenge for Latin America and the Caribbean Options for Climate-Resilient – Low-carbon Development*. Inter-American Development Bank; 2013.
 - The study deals with the impacts of climate change in LAC and how the consequences of climate change might affect the likelihood of achieving sustainable progress in the region. A substantial contribution of this report is the outlining of specific paths for mitigation and adaptation.
 2. United Nations Environmental Program (UNEP): *Vital Climate Change Graphics for Latin America and the Caribbean*. UNEP; 2010.
 3. Bouwman AF, Hoek KW, Van Der Olivier JGJ: **Uncertainties in the global source distribution of nitrous oxide inventories of N₂O**. *J Geophys Res* 1995, **100**:2785-2800.
 4. Austin AT, Bustamante MMC, Nardoto GB, Mitre SK, Perez T, Ometto JPHB, Ascarrunz NL, Forti MC, Longo K, Gavito ME, Enrich-Prast A, Martinelli LA: **Latin America's nitrogen challenge**. *Science* 2013, **340**:149.
 - The authors discuss the interrelated challenges in terms of human impact on the nitrogen (N) cycle with focus on Latin America. The main challenges associated with the N cycle and socio-economic issues of the region are presented.
 5. Ministerio del Ambiente y los Recursos Naturales (MARN): *Primera comunicación nacional em Cambio Climático de Venezuela*. [First National Communication on Climate Change of Venezuela]. Republica Bolivariana de Venezuela; 2005.
 6. Ministério da Ciência e Tecnologia e Inovação (MCTI): *Estimativas anuais de emissões de gases de efeito estufa no Brasil*. [Estimates of Annual Emissions of Greenhouse Gases in Brazil]. MCTI; 2013.
 7. Gobierno Argentino: *Segunda Comunicación Nacional del Gobierno Argentino a la Convención Marco de las Naciones Unidas sobre Cambio Climático*. [Second National Communication of the Argentine Government to the UN Framework Convention on Climate Change]. Proyecto BIRF No. TFO51287; 2007.
 8. SAGARPA-FAO: *Línea base del programa de sustentabilidad de los recursos naturales: Subíndice de emisiones de gases de efecto invernadero - metodología de cálculo*. [Baseline of the Program for Sustainability of Natural Resources: Sub-index of Emissions of Greenhouse Gases – Calculation Methodology]. Gobierno de México: Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación; 2012, .
 9. United Nations Environment Programme (UNEP): *Drawing Down N₂O to Protect Climate and the Ozone Layer. A UNEP Synthesis Report*. Nairobi, Kenya: United Nations Environment Programme (UNEP); 2013, .
 10. Martinelli LA: *Ecosystem Services and Agricultural Production in Latin America and Caribbean*. Technical Notes No. IDB-TN-382. Inter-American Development Bank; 2012.
 - The author presents a comprehensive analysis of the impact of agricultural development in LAC on biodiversity and associated ecosystem services.
 11. Olivier JGJ, Bouwman AF, van der Hoek KW, Berdowski JJM: **Global air emission inventories for anthropogenic sources of NO_x, NH₃ and N₂O in 1990**. *Environ Pollut* 1998, **102**:135-148.
 12. Bustamante MMC, Keller M, Silva DA: **Sources and sinks of trace gases in Amazonia and the Cerrado**. In *Amazonia and Global Change*. Edited by Keller M, Bustamante MMC, Gash J, Dias PL. American Geophysical Union; 2009:337-354.
 13. Aide TM, Clark ML, Grau HR, López-Carr D, Levy MA, Redo D, Bonilla-Moheno M, Riner G, Andrade-Núñez MJ, Muñiz M: **Deforestation and reforestation of Latin America and the Caribbean (2001–2010)**. *Biotropica* 2013, **45**:262-271.

This study evaluated deforestation and reforestation from the municipal to continental scale by producing annual maps of change in woody vegetation and other land-cover classes between 2001 and 2010 for each of the 16 050 municipalities in LAC. Furthermore, they determined which environmental or population variables best explained the variation in woody vegetation change.

14. Chuvieco E, Opazo S, Sione W, del Valle H, Anaya J, Di Bella C, Cruz I, Manzo L, López G, Mari N *et al.*: **Global burned-land estimation in Latin America using MODIS composite data.** *Ecol Appl* 2008, **18**:64-79.
15. Lauk C, Erb K-H: **Biomass consumed in anthropogenic vegetation fires: global patterns and processes.** *Ecol Econ* 2009, **69**:301-309.
16. Yevich R, Logan JA: **An assessment of biofuel use and burning of agricultural waste in the developing world.** *Glob Biogeochem Cycles* 2003, **17**:1095 <http://dx.doi.org/10.1029/2002GB001952>.
17. Longo K, Freitas SR, Andreae MO, Yokelson R, Artaxo P: **Biomass burning in Amazonia: emissions, long-range transport of smoke and its regional and remote impacts.** In *Amazonia and Global Change*. Edited by Keller M, Bustamante MMC, Gash J, Dias PL. American Geophysical Union; 2009.
18. De Vries JHM, Olivier JGJ, van den Wijngaart RA, Kreileman GJJ, Toer AMC: **Model for calculating regional energy use, industrial production and greenhouse gas emissions for evaluating global climate scenarios.** *Water Air Soil Pollut* 1994, **76**:79-131.
19. *Balanço Energético Nacional (BEN)*. [National Energy Balance]. Brasil: Ministério das Minas e Energia; 2013, .
20. Di Bella CM, Jobbágy EG, Paruelo JM, Pinnock S: **Continental fire density in South America.** *Glob Ecol Biogeogr* 2006, **15**:192-199.
21. Paruelo JM, Verón SR, Volante JN, Seghezzo L, Vallejos M, Aguiar S, Amdan L, Baldassini P, Ciuffolif L, Huykman N *et al.*: **Elementos conceptuales y metodológicos para la Evaluación de Impactos Ambientales Acumulativos (EIAAc) en bosques subtropicales: El caso del este de Salta, Argentina** [Conceptual and methodological elements for the Cumulative Environmental Impact Assessment (EIAAc) in subtropical forests: the case of eastern Salta, Argentina]. *Ecol Austral* 2011, **21**:163-178.
22. Grau HR, Aide TM, Gasparri NI: **Globalization and soybean expansion into semiarid ecosystems of Argentina.** *Ambio* 2005, **34**:265-266.
23. Grau HR, Gasparri NI, Aide TM: **Balancing food production and nature conservation in the neotropical dry forests of northern Argentina.** *Glob Change Biol* 2008, **14**:985-997.
24. CONAFOR: *Reporte semanal de resultados de incendios forestales*. [Weekly Report on Wildfires]. Coordinación General de Conservación y Restauración, Secretaría del Medio Ambiente y Recursos Naturales, Gobierno de México; 2013.
25. Berdanier AB, Conant RT: **Regionally differentiated estimates of cropland N₂O emissions reduce uncertainty in global calculations.** *Glob Change Biol* 2012, **18**:928-935.
26. Stehfest E, Bouwman L: **N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions.** *Nutr Cycl Agroecosyst* 2006, **74**:207-228.
27. Lapola DM, Martinelli LA, Peres CA, Ometto JPHB, Ferreira ME, Nobre CA, Aguiar APD, Bustamante MMC, Cardoso MF, Costa MH, Joly CA, Leite CC, Moutinho P, Sampaio G, Strassburg BBN, Vieira ICG: **Pervasive transition of the Brazilian land-use system.** *Nat Clim Change* 2013, **4**:27-35.
- The authors show that the relationship between agriculture, deforestation, and GHG emissions in Brazil has been changing, with the increasing intensification and conversion to commodity-based agriculture. While this is linked to the marked reduction in deforestation, these changes in the land-use system further stimulate social problems through increased inequality in land ownership and rural-urban migration.
28. INECC: *Inventario nacional de gases de efecto invernadero 1990-2010*. [National Inventory of Greenhouse Gases 1990-2010]. Secretaría del Medio Ambiente y Recursos Naturales, Gobierno de México; 2013.
29. García FM, Taboada MF, Gonzalez S, Picone L: *Alternativas para incrementar la eficiencia de uso en los cultivos de grano y mitigar las emisiones de óxido nítrico*. [Alternatives to Increase Use Efficiency in Grain Crops and Mitigate Nitrous Oxide Emissions]. Reporte Técnico IPNI/Fertilizar AC; 2013.
30. Marquina S, Donoso L, Pérez T, Gil J, Sanhueza E: **Losses of NO and N₂O emissions from Venezuelan and other worldwide tropical N-fertilized soils.** *J Geophys Res Biogeosci* 2013, **118** <http://dx.doi.org/10.1002/jgrg.20081>.
31. Crutzen PJ, Mosier AR, Smith KA, Winiwarter W: **N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuels.** *Atmos Chem Phys* 2008, **8**:389-395.
32. Carmo JB, Filoso S, Zotelli L, de Sousa Neto ER, Pitombo LM, Duarte-Neto PJ, Vargas VP, Andrade CA, Gava GC, Rossetto R, Cantarella H, Elia Neto A, Martinelli LA: **Infield greenhouse gas emissions from sugarcane soils in Brazil: effects from synthetic and organic fertilizer application and crop trash accumulation.** *GCB Bioenergy* 2012, **5**:1-14.
33. Vargas VP, Cantarella H, Martins AA, Soares JR, Carmo JB, Andrade CA: **Sugarcane crop residue increases N₂O and CO₂ emissions under high soil moisture conditions.** *Sugar Technol* 2014, **6**:174-179.
34. Austin AT, Piñeiro G, Gonzalez Polo M: **More is less: agricultural impacts on the N cycle in Argentina.** *Biogeochemistry* 2006, **79**:45-60.
35. Cruvinel EBF, Bustamante MMC, Kozovits AR, Zepp RG: **Soil emissions of NO, N₂O and CO₂ from croplands in the savanna region of central Brazil.** *Agric Ecosyst Environ* 2011, **144**:29-40.
36. Ciampitti IA, Ciarlo EA, Conti ME: **Nitrous oxide emissions from soil during soybean [*Glycine max* (L.) Merrill] crop phenological stages and stubbles decomposition period.** *Biol Fertil Soils* 2008, **44**:581-588.
37. Gerber PJ, Steinfeld H, Henderson B, Mottet A, Opio C, Dijkman J, Falucco A, Tempio G: *Tackling Climate Change Through Livestock – A Global Assessment of Emissions and Mitigation Opportunities*. Rome: Food and Agriculture Organization of the United Nations (FAO); 2013, .
38. Lessa ACR, Madari BE, Paredes DS, Boddey RM, Urquiaga S, Jantalia CP, Alves BJR: **Bovine urine and dung deposited on Brazilian savannah pastures contribute differently to direct and indirect soil nitrous oxide emissions.** *Agric Ecosyst Environ* 2014 <http://dx.doi.org/10.1016/j.agee.2014.01.010>.
39. FRA: *Global Forest Resources Assessment, FAO Forestry Paper 163*. Rome, Italy: FAO; 2010, .
40. Cole CV, Duxbury J, Freney J, Heinemeyer O, Minami K, Mosier A, Paustin K, Rosenberg N, Sampson N, Sauerbeck D *et al.*: **Global estimates of potential mitigation of greenhouse gas emissions by agriculture.** *Nutr Cycl Agroecosyst* 1997, **49**:221-228.
41. Signor D, Cerri CEP, Conant R: **N₂O emissions due to nitrogen fertilizer applications in two regions of sugarcane cultivation in Brazil.** *Environ Res Lett* 2013, **8**:015013.
42. Subbarao GV, Nakahara K, Hurtado MP, Ono H, Moreta DE, Salcedo AF, Yoshihashi AT, Ishikawa T, Ishitani M, Ohnishi-Kameyama M *et al.*: **Evidence for biological nitrification inhibition in *Brachiaria* pastures.** *Proc Natl Acad Sci U S A* 2009, **106**:17302-17307.
43. Ruan L, Robertson GP: **Initial nitrous oxide, carbon dioxide, and methane costs of converting conservation reserve program grassland to row crops under no-till vs. conventional tillage.** *Glob Change Biol* 2013, **19**:2478-2489.
44. Van Kessel C, Venterea R, Six J, Adviento-Borbe MA, Linquist B, van Groenigen KJ: **Climate, duration, and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis.** *Glob Change Biol* 2013, **19**:33-44.
45. Boddey RM, Jantalia CP, Conceição PC, Zanata JA, Bayer C, Mielniczuk J, Dieckow J, Santos HP, dos Denardin JE, Aita C *et al.*: **Carbon accumulation at depth in Ferralsols under zero-till subtropical agriculture.** *Glob Change Biol* 2010, **16**:784-795.
46. Viglizzo EF, Frank CF, Carreño LV, Jobbágy EG, Pereyra H, Clatt J, Pincén D, Ricard MF: **Ecological and environmental footprint of 50 years of agricultural expansion in Argentina.** *Glob Change Biol* 2011, **17**:959-973.

47. Astier M, Merlín-Uribe Y, Villamil-Echeverri L, Garcíarreal A, Gavito ME, Masera OR: **Energy balance and greenhouse gas emissions in organic and conventional avocado orchards in Mexico.** *Ecol Indic* 2014, **43**:281-287.
48. Skinner C, Gattinger A, Muller A, Mäder P, Fliessbach A, Stolze M, Ruser R, Niggli U: **Greenhouse gas fluxes from agricultural soils under organic and non-organic management — a global meta-analysis.** *Sci Total Environ* 2014, **468–469**:553-563.
49. Bouwman G, Van Drecht G, van der Hoek KW: **Surface N balances and reactive N loss to the environment from global intensive agricultural production systems for the period 1970–2030.** *Sci China C Life Sci* 2005, **48(Suppl 1–13)**:1.