

Letters

Photoreceptors UVR8 and phytochrome B cooperate to optimize plant growth and defense in patchy canopies

Light is a critical source of information for plants. Plants use the phytochromes (particularly phyB) to detect light signals associated with the proximity of competitors. A low ratio of red (R) to far-red (FR) radiation (R : FR) indicates increased competition intensity, and triggers morphological responses that allow the plant to escape shading from its neighbors (the shade avoidance syndrome, SAS) (reviewed in Ballaré, 2009; Kami *et al.*, 2010; Martínez-García *et al.*, 2010; Casal, 2012; Pierik & de Wit, 2014). Recent evidence from studies on light regulation of plant immunity has suggested that plants may also use ultraviolet-B (UV-B, 290–315 nm) radiation as an indicator of competition intensity and light availability (Demkura *et al.*, 2010; Demkura & Ballaré, 2012). In addition, recent studies have shown that UV-B radiation can strongly repress SAS responses triggered by low R : FR ratios (Hayes *et al.*, 2014). Ambient UV-B radiation causes damaging effects on plants, such as DNA damage (Mazza *et al.*, 1999), and also induces adaptive photomorphogenic responses acting through a specific UV-B photoreceptor (UVR8) (Rizzini *et al.*, 2011; Christie *et al.*, 2012; Wu *et al.*, 2012) (reviewed in Heijde & Ulm, 2012; Jenkins, 2014). Therefore, the possibility exists that plants integrate information perceived by phyB and UVR8 to make decisions about growth and defense when faced with a complex light environment, such as the one that characterizes vegetation canopies. In this Letter, we address this possibility and discuss how the interplay between UV-B and R : FR signaling fine tunes plant growth and defense to optimize resource utilization in patchy canopy environments.

UV-B tempers SAS responses triggered by phyB inactivation under sunlight

In *Arabidopsis* seedlings grown under artificial light, moderate levels of supplemental UV-B can strongly suppress hypocotyl elongation responses to low R : FR ratio (Hayes *et al.*, 2014). However, under field conditions, phyB inactivation can trigger significant elongation responses even in plants growing under full sunlight. Perhaps the most dramatic example comes from studies with cucumber (*Cucumis sativus*), which showed that the phyB null mutant (*lh*) can grow hypocotyls as long as 30 cm even under summertime field conditions and high natural UV-B irradiance (Casal *et al.*, 1994). To further explore the interactions between phyB inactivation and natural UV-

B levels, we compared the growth of *lh* and wild type cucumbers under filters that either blocked (–UV-B) or transmitted UV-B radiation (+UV-B) (see Supporting Information Methods S1). We found a dramatic effect of phyB inactivation promoting hypocotyl elongation and a modest, but significant repression of elongation by solar UV-B radiation (Fig. 1a). Less spectacular responses to partial phyB inactivation have been reported in other field studies, under solar UV-B irradiances, using other plant species exposed to lateral FR radiation (reviewed in Ballaré, 1999). Since none of these field studies involved *Arabidopsis* plants, we characterized SAS responses to the proximity of competitors in wild-type and *uvr8* *Arabidopsis* seedlings grown in the field under contrasting levels of UV-B radiation (Fig. 1b). Plants responded to the proximity of grass competitors with a rapid increase in leaf angles (hyponastic response). This hyponastic response was reduced by solar UV-B radiation, and the repressive effect of solar UV-B radiation was missing in a *uvr8* mutant (Fig. 1c). Exploration of *Arabidopsis* transcriptome data in microarray databases (Tables S1–S3) further suggest the existence of antagonistic effects of low R : FR ratio and UV-B radiation on expression of growth (auxin)-related genes (Fig. 2a). For example *YUCCA9*, *ATHB-2*, *IAA29* and *SAUR23* were all strongly upregulated by low R : FR and downregulated by UV-B radiation (Table S3). Significant effects of UVR8 activation reducing the expression of selected auxin-related genes have been reported recently (Vandenbussche *et al.*, 2014), particularly in plants exposed to low R : FR ratios (including *YUCCA9* and *IAA29*; Hayes *et al.*, 2014). In our meta-analysis of available microarray data, the effect of UV-B on the ‘Auxin’ gene ontology (GO) category appeared to be somewhat dependent on the experimental conditions (Fig. S1) and largely conserved in *uvr8* (Fig. S2). Taken together, physiological (Fig. 1) and transcriptomic (Fig. 2a; Table S3) results are consistent with the idea that solar UV-B radiation moderates the growth responses to neighbor proximity triggered by low R : FR ratios. The magnitude of the effect of UV-B (and UVR8) is likely to vary with the relative levels of R : FR and UV-B radiation, which emphasizes the need for using experimental approaches that cover the natural range of variation of these light signals.

UV-B and high R : FR promote plant defense under sunlight, via partially overlapping mechanisms

Solar UV-B radiation promotes plant defense against herbivorous insects and some pathogens, and here again the effects of solar UV-B are opposite to those of low R : FR ratio (Ballaré *et al.*, 2012). The positive effect of (natural) UV-B radiation on plant immunity has often been attributed to the accumulation of leaf phenolics, although enhanced signaling through the jasmonate (JA) defense pathway is also thought to play an important role (Ballaré, 2014).




Fig. 1 The interplay between solar ultraviolet-B (UV-B) and phyB in shaping plant morphology under field conditions. (a) phyB inactivation in the *lh* mutant (López-Juez *et al.*, 1992) triggers a strong shade avoidance response in field grown plants, and solar UV-B represses elongation. Average hypocotyl length (cm) for wild-type (WT) plants was 5.9 (+UV-B) and 7.1 (-UV-B), and for *lh* plants 10.9 (+UV-B) and 13.0 (-UV-B); *lh* effect $P < 0.01$; UV-B effect $P = 0.05$; interaction $P = 0.56$; $n = 5$; see Supporting information Methods S1 for experimental details; filter types and configuration were as described previously (Izaguirre *et al.*, 2007; Zavala *et al.*, 2014). (b) Experimental setup used to test Arabidopsis responses to the proximity of grass neighbors in the field under contrasting levels of solar UV-B radiation. Wild-type (Col-0) and *uvr8-6* mutant (Favory *et al.*, 2009) Arabidopsis plants were surrounded by zero (D0), three (D3), or six (D6) ryegrass-containing pots (only treatments D6 (top, six ryegrass pots) and D3 (bottom, three ryegrass pots and three pots containing only soil) are shown in the picture); the setup was replicated four times under either solar UV-B (+UV-B) or attenuated UV-B (-UV-B); see Methods S1 for details. (c) Arabidopsis plants respond to the proximity of grass neighbors by increasing leaf angles, and ambient UV-B tempers this response via UVR8. Bars, ± 1 standard error (SE).

Leaf phenolics

Accumulation of soluble phenolic compounds is one of the most ubiquitous responses to UV-B radiation (Searles *et al.*, 2001), which is to be expected given the key role played by these compounds in UV-photoprotection (Braun & Tevini, 1993; Mazza *et al.*, 2000). In Arabidopsis, the accumulation of both phenylpropanoids (sinapates) and flavonoids is boosted by UV-B radiation (Li *et al.*, 1993; Landry *et al.*, 1995), in a UVR8-dependent manner (Kliebenstein *et al.*, 2002; Demkura & Ballaré,




Fig. 2 Comparison of the effects of low red : far red (R : FR) and supplemental ultraviolet-B (UV-B) radiation on gene expression for genes belonging to selected gene ontology (GO) categories. Data are from public databases (see Supporting Information Methods S1 for details) and many of them have been reported in previous publications (Brown *et al.*, 2005; Tao *et al.*, 2008; Favory *et al.*, 2009; Kozuka *et al.*, 2010; Cerrudo *et al.*, 2012; Leivar *et al.*, 2012; De Wit *et al.*, 2013; Morales *et al.*, 2013; Reddy *et al.*, 2013). Data points highlighted in red (a) correspond to *YUCCA9* (At1G04180), *ATHB-2* (At4 g16780), *IAA29* (At4 g32280) and *SAUR23* (At5 g18060); see Table S3 for quantitative details.

2012). Some of these compounds (sinapates) have been shown to mediate the positive effect of UV-B radiation on plant immunity against fungal pathogens (Demkura & Ballaré, 2012). In contrast to the consistent effect of UV-B, the effects of R : FR ratio on levels of soluble leaf phenolics can be quite variable (Table 1), and often not significant. These observations at the metabolite level are totally

consistent with our meta-analysis of transcriptome data in *Arabidopsis*: supplemental UV-B has a robust effect increasing the expression of phenylpropanoid- and flavonoid-related genes, whereas the effects of FR are generally very modest (Fig. 2b,c; Table S3). The effects of UV-B on phenylpropanoid- and flavonoid-related genes are mostly mediated by UVR8 (Fig. S2b,c).

JA signaling

Reductions in UV-B radiation and R:FR ratio may have convergent effects repressing JA signaling. Genetic evidence in *Arabidopsis* and *Nicotiana* indicates that some of the anti-herbivore effects of solar UV-B are missing in plants impaired in the biosynthesis of bioactive JA (Caputo *et al.*, 2006; Demkura *et al.*, 2010). Solar UV-B can increase JA synthesis or expression of JA biosynthetic genes in *Nicotiana* spp. (Izaguirre *et al.*, 2003; Dinh *et al.*, 2012), although this is not always the case (Demkura *et al.*, 2010). In species of the Solanaceae, UV-B increases plant sensitivity to JA for activation of proteinase inhibitors (Stratmann *et al.*, 2000; Demkura *et al.*, 2010). However, this enhancing effect was not detected in *Arabidopsis* for other markers of the JA response (Demkura & Ballaré, 2012). Available microarray data indicate that UV-B radiation increases the expression of some JA-related genes (Fig. 2d; see also Table S3 and Fig. S1), but this effect appears to be highly dependent on the experimental conditions (Fig. S1xiii–xvi), and to some extent conserved in *uvr8* mutants (Fig. S2d). The effects of low R:FR reducing JA sensitivity are very well documented (reviewed in Ballaré, 2014). Transcriptome data in *Arabidopsis* do not reveal major effects of R:FR ratio on JA-related genes in *healthy* (non-induced) plants (Fig. 2d), which is consistent with the idea that R:FR regulates JA signaling predominantly by reducing the response to the JA burst produced by the plant in response to herbivory or pathogen attack (Moreno *et al.*, 2009;

Cerrudo *et al.*, 2012; De Wit *et al.*, 2013). The effect of low R:FR ratios repressing the induction of JA-dependent genes correlates with the down-regulation of specific metabolites involved in direct (Cargnel *et al.*, 2014) and indirect (Izaguirre *et al.*, 2013; Kegge *et al.*, 2013) defenses. From an ecological point of view, down-regulation of JA responses under conditions of low R:FR and UV-B irradiance is likely to help the plant to redirect resources from defense to rapid elongation, thereby increasing its ability to compete for light.

Interactions between UVR8 and phyB signaling mediated by hormonal cross-talk

Recent work on the effects of R:FR and UV-B on growth- and defense-related hormonal pathways (Hayes *et al.*, 2014; Leone *et al.*, 2014) can provide important clues to understand the mechanisms by which R:FR and UV-B signals are used by the plant to optimize resource allocation in patchy canopies (Fig. 3). Reduced JA signaling under low R:FR ratios has been attributed to increased stability of JAZ10 (Leone *et al.*, 2014) (and presumably other JAZ proteins – Chico *et al.*, 2014), and increased turnover of DELLA proteins (Leone *et al.*, 2014). JAZs are key repressors of JA signaling (Browse, 2009; Fonseca *et al.*, 2009; Kazan & Manners, 2012), and DELLA proteins, which are key repressors of gibberellin (GA) signaling (Harberd *et al.*, 2009), are known to recruit JAZs into inactive protein complexes (Hou *et al.*, 2010; Yang *et al.*, 2012). Hayes *et al.* (2014) reported that UV-B supplementation can stabilize DELLAs. Therefore, it is conceivable that changes in DELLA turnover are functionally important not only to define the balance between UV-B and low R:FR in the modulation of growth responses, but also in the regulation of JA signaling and plant immunity in canopies (Fig. 3).

Table 1 Far red (FR) provided against a background of white light to lower the red (R) to far red (R:FR) ratio has mixed effects on accumulation of soluble phenolic compounds in the leaves of de-etiolated plants

Species	Effect of supplemental FR (low R:FR ratio)	Effect of supplemental UV-B under comparable physiological conditions ¹
<i>Betula pendula</i>	No effect on flavonoids; marginal increase in the contents of phenolic acids, such as chlorogenic acid and cinnamic acid (Tegelberg <i>et al.</i> , 2004)	Increased flavonoids and phenolic acids (Tegelberg <i>et al.</i> , 2004)
<i>Nicotiana longiflora</i>	Reduced the accumulation of herbivory-induced chlorogenic acid (Izaguirre <i>et al.</i> , 2006)	Increased flavonoids and phenolic acids (Izaguirre <i>et al.</i> , 2007)
<i>Solanum lycopersicum</i>	Increased accumulation of flavonoids in the leaves (the opposite effect was observed for stem flavonoids and anthocyanins) (Cagnola <i>et al.</i> , 2012)	Increased flavonoid accumulation (Ballaré <i>et al.</i> , 1995)
<i>Arabidopsis thaliana</i>	No effect on leaf flavonoids or sinapates (Cargnel <i>et al.</i> , 2014)	Increased flavonoids and sinapates in an UVR8-dependent manner (Demkura & Ballaré, 2012)
<i>Arabidopsis thaliana</i>	Decreased content of anthocyanins in plants treated with methyl-jasmonate (Cerrudo <i>et al.</i> , 2012; Leone <i>et al.</i> , 2014)	Not reported

¹Results are shown for comparison. In all the UV-B supplementation experiments listed in this column, plants were exposed to physiological levels of UV-B radiation against a background of white light.




Fig. 3 UVR8 and phyB cooperatively modulate plant phenotypic plasticity in plant canopies. Hypothetical mechanism by which changes in red : far-red (R : FR), perceived by phyB, and UV-B radiation, perceived by UVR8, regulate growth and defense in response to shading and neighbor proximity. (a) In canopy gaps and when neighbors are far away, phyB Pfr levels are high, which reduce the levels and activity of PHYTOCHROME INTERACTING FACTORS (PIFs) (Lorrain *et al.*, 2008; Hornitschek *et al.*, 2012; Li *et al.*, 2012; Park *et al.*, 2012). PIFs are also repressed by abundant DELLA proteins (Feng *et al.*, 2008; de Lucas *et al.*, 2008), which also keep JASMONATE ZIM domain (JAZ) proteins from repressing their target transcription factors, such as MYCs (Hou *et al.*, 2010). At the same time, activation of UVR8 by solar UV-B facilitates accumulation of DELLA proteins in a HY5-dependent manner (likely by increasing gibberellin (GA) degradation), and promotes PIF degradation in a HY5-independent manner (Hayes *et al.*, 2014). Under these conditions, auxin biosynthesis and elongation growth are repressed, and jasmonate (JA)-dependent defense responses can be readily activated following JA-induced JAZ protein degradation. UV-B, acting through UVR8, promotes the accumulation of soluble phenolic compounds (flavonoids and phenylpropanoids), which can contribute to increase plant defense in a JA-independent manner (Demkura & Ballaré, 2012). (b) Under shade or high density conditions, low R : FR ratios lead to phyB inactivation, increased levels and activity of PIFs, increased auxin biosynthesis (Tao *et al.*, 2008), and rapid degradation of DELLAs (Leone *et al.*, 2014), thereby promoting elongation. Low UV-B levels also reduce UVR8-induced PIF turnover and facilitate GA accumulation and degradation of DELLA proteins (Hayes *et al.*, 2014). DELLA degradation frees up JAZ proteins, which are also stabilized (Chico *et al.*, 2014; Leone *et al.*, 2014) and are therefore present and available to repress defense-activating transcription factors. Under these conditions, MYC levels can also be reduced (Chico *et al.*, 2014), which further suppresses defense responses. Arrows, promotion; blunt lines, repression/inactivation. Line thickness indicates the strength of the effect; the relative size of the ovals denotes concentration or activity, and overlap indicates an antagonistic interaction between proteins. Dashed arrows, inducible responses. JA-Ile, jasmonyl-L-isoleucine; TF, transcription factor.

Concluding remarks

UVR8 and phyB cooperatively modulate the balance between growth and defense through both convergent and parallel signaling pathways (Fig. 3). The use of common signaling elements (such as DELLA proteins), provides a simple mechanism to reinforce the response when both photoreceptors sense contingent information (e.g. under deep canopy shade or in the open), or to balance out their effects when they perceive conflicting canopy signals (e.g. during proximity perception). In the latter case, FR reflected by neighboring plants provides the plant with a key warning signal of oncoming competition when light levels are still high (Ballaré *et al.*, 1990). However, because excessive responses to low R : FR may entail fitness costs (Casal *et al.*, 1994; Dorn *et al.*, 2000; McGuire & Agrawal, 2005), moderation of these responses by UVR8 in canopy gaps (Fig. 1) may be ecologically advantageous. In addition, the use of independent signaling pathways may allow for one photoreceptor to dominate the pattern of response. This appears to be the case for accumulation of phenolic sunscreens, where UVR8 has a distinctly dominant role (Table 1; Fig. 2). Since UV-B can penetrate more in leaf canopies than longer wavelengths (Flint & Caldwell, 1998), an overriding role of the UVR8 pathway may induce adaptive UV-B photoprotection even under conditions of low R : FR. The emerging map of interactions between UVR8 and

phyB signaling (Fig. 3) is revealing exciting new insights into the mechanisms used by plants to optimize growth and defense plasticity in variable light environments.

Acknowledgements

This research was financially supported by grants from CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas), ANPCyT (Agencia Nacional de Promoción Científica y Tecnológica) and UBACyT (Universidad de Buenos Aires Ciencia y Técnica). The authors thank Amy Austin for helpful discussions and comments on the manuscript and Pablo Rosano for his assistance in the field experiments.

Carlos A. Mazza¹ and Carlos L. Ballaré^{1,2*}

¹IFEVA, Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad de Buenos Aires, C1417DSE Buenos Aires, Argentina;

²IIB-INTECH, Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad Nacional de San Martín, B1650HMP Buenos Aires, Argentina

(*Author for correspondence: tel +54 11 4524 8070; email ballare@ifeva.edu.ar)

References

- Ballaré CL. 1999. Keeping up with the neighbours: phytochrome sensing and other signalling mechanisms. *Trends in Plant Science* 4: 97–102.
- Ballaré CL. 2009. Illuminated behaviour: phytochrome as a key regulator of light foraging and plant anti-herbivore defence. *Plant, Cell & Environment* 32: 713–725.
- Ballaré CL. 2014. Light regulation of plant defense. *Annual Review of Plant Biology* 65: 335–363.
- Ballaré CL, Barnes PW, Flint SD, Price S. 1995. Inhibition of hypocotyl elongation by ultraviolet-B radiation in de-etiolating tomato seedlings. II. Time-course, comparison with flavonoid responses and adaptive significance. *Physiologia Plantarum* 93: 593–601.
- Ballaré CL, Mazza CA, Austin AT, Pierik R. 2012. Canopy light and plant health. *Plant Physiology* 160: 145–155.
- Ballaré CL, Scopel AL, Sánchez RA. 1990. Far-red radiation reflected from adjacent leaves: an early signal of competition in plant canopies. *Science* 247: 329–332.
- Braun J, Tevini M. 1993. Regulation of UV-protective pigment synthesis in the epidermal layer of rye seedlings (*Secale cereale* L. cv. Kustro). *Photochemistry and Photobiology* 57: 318–323.
- Brown BA, Cloix C, Jiang GH, Kaiserli E, Herzyk P, Kliebenstein DJ, Jenkins GI. 2005. A UV-B-specific signaling component orchestrates plant UV protection. *Proceedings of the National Academy of Sciences, USA* 102: 18225–18230.
- Browse J. 2009. Jasmonate passes muster: a receptor and targets for the defense hormone. *Annual Review of Plant Biology* 60: 183–205.
- Cagnola JJ, Ploschuk E, Benech-Arnold T, Finlayson SA, Casal JJ. 2012. Stem transcriptome reveals mechanisms to reduce the energetic cost of shade-avoidance responses in tomato. *Plant Physiology* 160: 1110–1119.
- Caputo C, Rutitzky M, Ballaré CL. 2006. Solar ultraviolet-B radiation alters the attractiveness of *Arabidopsis* plants to diamondback moths (*Plutella xylostella* L.): impacts on oviposition and involvement of the jasmonic acid pathway. *Oecologia* 149: 81–90.
- Cargnel MD, Demkura PV, Ballaré CL. 2014. Linking phytochrome to plant immunity: low red : far-red ratios increase *Arabidopsis* susceptibility to *Botrytis cinerea* by reducing the biosynthesis of indolic glucosinolates and camalexin. *New Phytologist* 204: 342–354.
- Casal JJ. 2012. Shade avoidance. *Arabidopsis Book* 10: e0157.
- Casal JJ, Ballaré CL, Tourn M, Sanchez RA. 1994. Anatomy, growth and survival of a long-hypocotyl mutant of *Cucumis sativus* deficient in phytochrome b. *Annals of Botany* 73: 569–575.
- Cerrudo I, Keller MM, Cargnel MD, Demkura PV, de Wit M, Patitucci MS, Pierik R, Pieterse CMJ, Ballaré CL. 2012. Low red/far-red ratios reduce *Arabidopsis* resistance to *Botrytis cinerea* and jasmonate responses via a COI1-JAZ10-dependent, salicylic acid-independent mechanism. *Plant Physiology* 158: 2042–2052.
- Chico J-M, Fernández-Barbero G, Chini A, Fernández-Calvo P, Díez-Díaz M, Solano R. 2014. Repression of jasmonate-dependent defenses by shade involves differential regulation of protein stability of MYC transcription factors and their JAZ repressors in *Arabidopsis*. *Plant Cell* 26: 1967–1980.
- Christie JM, Arvai AS, Baxter KJ, Heilmann M, Pratt AJ, O'Hara A, Kelly SM, Hothorn M, Smith BO, Hitomi K *et al.* 2012. Plant UVR8 photoreceptor senses UV-B by tryptophan-mediated disruption of cross-dimer salt bridges. *Science* 335: 1492–1496.
- De Wit M, Spoel SH, Sanchez Perez GF, Gommers CMM, Pieterse CMJ, Voisenek LACJ, Pierik R. 2013. Perception of low red:far-red ratio compromises both salicylic acid- and jasmonic acid-dependent pathogen defences in *Arabidopsis*. *Plant Journal* 75: 90–103.
- Demkura PV, Abdala G, Baldwin IT, Ballaré CL. 2010. Jasmonate-dependent and -independent pathways mediate specific effects of solar ultraviolet-B radiation on leaf phenolics and antiherbivore defense. *Plant Physiology* 152: 1084–1095.
- Demkura PV, Ballaré CL. 2012. UVR8 mediates UV-B-induced *Arabidopsis* defense responses against *Botrytis cinerea* by controlling sinapate accumulation. *Molecular Plant* 5: 642–652.
- Dinh ST, Gális I, Baldwin IT. 2012. UVB radiation and HGL-DTGs provide durable resistance against mirid (*Tupiocoris notatus*) attack in field-grown *Nicotiana attenuata* plants. *Plant, Cell & Environment* 36: 590–606.
- Dorn LA, Pyle EH, Schmitt J. 2000. Plasticity to light cues and resources in *Arabidopsis thaliana*: testing for adaptive value and costs. *Evolution* 54: 1982–1994.
- Favory JJ, Stec A, Gruber H, Rizzini L, Oravec A, Funk M, Albert A, Cloix C, Jenkins GI, Oakeley EJ *et al.* 2009. Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress acclimation in *Arabidopsis*. *EMBO Journal* 28: 591–601.
- Feng SH, Martinez C, Gusmaroli G, Wang Y, Zhou JL, Wang F, Chen LY, Yu L, Iglesias-Pedraz JM, Kircher S *et al.* 2008. Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* 451: 475–479.
- Flint SD, Caldwell MM. 1998. Solar UV-B and visible radiation in tropical forest gaps: measurements partitioning direct and diffuse radiation. *Global Change Biology* 4: 863–870.
- Fonseca S, Chico JM, Solano R. 2009. The jasmonate pathway: the ligand, the receptor and the core signalling module. *Current Opinion in Plant Biology* 12: 539–547.
- Harberd NP, Belfield E, Yasumura Y. 2009. The angiosperm gibberellin-GID1-DELLA growth regulatory mechanism: how an “inhibitor of an inhibitor” enables flexible response to fluctuating environments. *Plant Cell* 21: 1328–1339.
- Hayes S, Velanis CN, Jenkins GI, Franklin KA. 2014. UV-B detected by the UVR8 photoreceptor antagonizes auxin signaling and plant shade avoidance. *Proceedings of the National Academy of Sciences, USA* 111: 11894–11899.
- Heijde M, Ulm R. 2012. UV-B photoreceptor-mediated signalling in plants. *Trends in Plant Science* 17: 230–237.
- Hornitschek P, Kohnen MV, Lorrain S, Rougemont J, Ljung K, López-Vidriero I, Franco-Zorrilla JM, Solano R, Trevisan M, Pradervand S *et al.* 2012. Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling. *Plant Journal* 71: 699–711.
- Hou X, Lee LYC, Xia K, Yan Y, Yu H. 2010. DELLAs modulate jasmonate signaling via competitive binding to JAZs. *Developmental Cell* 19: 884–894.
- Izaguirre MM, Mazza CA, Astigueta MS, Ciarla AM, Ballaré CL. 2013. No time for candy: passionfruit (*Passiflora edulis*) plants down-regulate damage-induced extra floral nectar production in response to light signals of competition. *Oecologia* 173: 213–221.
- Izaguirre MM, Mazza CA, Biondini M, Baldwin IT, Ballaré CL. 2006. Remote sensing of future competitors: impacts on plant defenses. *Proceedings of the National Academy of Sciences, USA* 103: 7170–7174.
- Izaguirre MM, Mazza CA, Svatos A, Baldwin IT, Ballaré CL. 2007. Solar ultraviolet-B radiation and insect herbivory trigger partially overlapping phenolic responses in *Nicotiana attenuata* and *Nicotiana longiflora*. *Annals of Botany* 99: 103–109.
- Izaguirre MM, Scopel AL, Baldwin IT, Ballaré CL. 2003. Convergent responses to stress. Solar ultraviolet-B radiation and *Manduca sexta* herbivory elicit overlapping transcriptional responses in field-grown plants of *Nicotiana longiflora*. *Plant Physiology* 132: 1755–1767.
- Jenkins GI. 2014. The UV-B photoreceptor UVR8: from structure to physiology. *Plant Cell* 26: 21–37.
- Kami C, Lorrain S, Hornitschek P, Fankhauser C. 2010. Light-regulated plant growth and development. In: Timmermans MCP, ed. *Current topics in developmental biology*. New York, NY, USA: Academic Press, 29–66.
- Kazan K, Manners JM. 2012. JAZ repressors and the orchestration of phytohormone crosstalk. *Trends in Plant Science* 17: 22–31.
- Kegge W, Weldegergis BT, Soler R, Eijk MV-V, Dicke M, Voisenek LACJ, Pierik R. 2013. Canopy light cues affect emission of constitutive and methyl jasmonate-induced volatile organic compounds in *Arabidopsis thaliana*. *New Phytologist* 200: 861–874.
- Kliebenstein DJ, Lim JE, Landry LG, Last RL. 2002. *Arabidopsis* UVR8 regulates ultraviolet-B signal transduction and tolerance and contains sequence similarity to human regulator of chromatin condensation. *Plant Physiology* 130: 234–243.
- Kozuka T, Kobayashi J, Horiguchi G, Demura T, Sakakibara H, Tsukaya H, Nagatani A. 2010. Involvement of auxin and brassinosteroid in the regulation of petiole elongation under the shade. *Plant Physiology* 153: 1608–1618.

- Landry LG, Chapple CC, Last RL. 1995. *Arabidopsis* mutants lacking phenolic sunscreens exhibit enhanced ultraviolet-B injury and oxidative damage. *Plant Physiology* 109: 1159–1166.
- Leivar P, Tepperman JM, Cohn MM, Monte E, Al-Sady B, Erickson E, Quail PH. 2012. Dynamic antagonism between phytochromes and PIF family basic helix-loop-helix factors induces selective reciprocal responses to light and shade in a rapidly responsive transcriptional network in *Arabidopsis*. *Plant Cell* 24: 1398–1419.
- Leone M, Keller MM, Cerrudo I, Ballaré CL. 2014. To grow or defend? Low red:far-red ratios reduce jasmonate sensitivity in *Arabidopsis* seedlings by promoting DELLA degradation and increasing JAZ10 stability. *New Phytologist* 204: 355–367.
- Li J, Ou-Lee TM, Raba R, Amundson RG, Last RL. 1993. *Arabidopsis* flavonoid mutants are hypersensitive to UV-B irradiation. *Plant Cell* 5: 171–179.
- Li L, Ljung K, Breton G, Schmitz RJ, Pruneda-Paz J, Cowing-Zitron C, Cole BJ, Ivans LJ, Pedmale UV, Jung H-S *et al.* 2012. Linking photoreceptor excitation to changes in plant architecture. *Genes & Development* 26: 785–790.
- López-Juez E, Nagatani A, Tomizawa KI, Deak M, Kern R, Kendrick RE, Furuya M. 1992. The cucumber long hypocotyl mutant lacks a light-stable PHYB-like phytochrome. *Plant Cell* 4: 241–251.
- Lorrain S, Allen T, Duek PD, Whitelam GC, Fankhauser C. 2008. Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant Journal* 53: 312–323.
- de Lucas M, Daviere JM, Rodriguez-Falcon M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blazquez MA, Titarenko E, Prat S. 2008. A molecular framework for light and gibberellin control of cell elongation. *Nature* 451: 480–484.
- Martínez-García JF, Galstyan A, Salla-Martret M, Cifuentes-Esquivel N, Gallemí M, Bou-Torrent J, Jean-Claude Kader, Michel D. 2010. Regulatory components of shade avoidance syndrome. *Advances in Botanical Research* 53: 65–116.
- Mazza CA, Battista D, Zima AM, Szwarcberg-Bracchitta M, Giordano CV, Acevedo A, Scopel AL, Ballaré CL. 1999. The effects of solar ultraviolet-B radiation on the growth and yield of barley are accompanied by increased DNA damage and antioxidant responses. *Plant, Cell & Environment* 22: 61–70.
- Mazza CA, Boccalandro HE, Giordano CV, Battista D, Scopel AL, Ballaré CL. 2000. Functional significance and induction by solar radiation of ultraviolet-absorbing sunscreens in field-grown soybean crops. *Plant Physiology* 122: 117–125.
- McGuire R, Agrawal AA. 2005. Trade-offs between the shade-avoidance response and plant resistance to herbivores? Tests with mutant *Cucumis sativus*. *Functional Ecology* 19: 1025–1031.
- Morales LO, Brosché M, Vainonen J, Jenkins GI, Wargent JJ, Sipari N, Strid A, Lindfors AV, Tegelberg R, Aphalo PJ. 2013. Multiple roles for UV RESISTANCE LOCUS8 in regulating gene expression and metabolite accumulation in *Arabidopsis* under solar ultraviolet radiation. *Plant Physiology* 161: 744–759.
- Moreno JE, Tao Y, Chory J, Ballaré CL. 2009. Ecological modulation of plant defense via phytochrome control of jasmonate sensitivity. *Proceedings of the National Academy of Sciences, USA* 106: 4935–4940.
- Park E, Park J, Kim J, Nagatani A, Lagarias JC, Choi G. 2012. Phytochrome B inhibits binding of phytochrome-interacting factors to their target promoters. *Plant Journal* 72: 537–546.
- Pierik R, de Wit M. 2014. Shade avoidance: phytochrome signalling and other aboveground neighbour detection cues. *Journal of Experimental Botany* 65: 2815–2824.
- Reddy SK, Holalu SV, Casal JJ, Finlayson SA. 2013. Abscisic acid regulates axillary bud outgrowth responses to the ratio of red to far-red light. *Plant Physiology* 163: 1047–1058.
- Rizzini L, Favory JJ, Cloix C, Faggionato D, O'Hara A, Kaiserli E, Baumeister R, Schäfer E, Nagy F, Jenkins GI *et al.* 2011. Perception of UV-B by the *Arabidopsis* UVR8 protein. *Science* 332: 103–106.
- Searles PS, Flint SD, Caldwell MM. 2001. A meta analysis of plant field studies simulating stratospheric ozone depletion. *Oecologia* 127: 1–10.
- Stratmann JW, Stelmach BA, Weiler EW, Ryan CA. 2000. UVB/UVA radiation activates a 48 kDa myelin basic protein kinase and potentiates wound signaling in tomato leaves. *Photochemistry and Photobiology* 71: 116–123.
- Tao Y, Ferrer JL, Ljung K, Pojer F, Hong F, Long JA, Li L, Moreno JE, Bowman ME, Ivans LJ *et al.* 2008. Rapid synthesis of auxin via a new tryptophan-dependent pathway is required for shade avoidance in plants. *Cell* 133: 164–176.
- Tegelberg R, Julkunen-Tiitto R, Aphalo PJ. 2004. Red:far-red light ratio and UV-B radiation: their effects on leaf phenolics and growth of silver birch seedlings. *Plant, Cell & Environment* 27: 1005–1013.
- Vandenbussche F, Tilbrook K, Fierro AC, Marchal K, Poelman D, Van Der Straeten D, Ulm R. 2014. Photoreceptor-mediated bending towards UV-B in *Arabidopsis*. *Molecular Plant* 7: 1041–1052.
- Wu D, Hu Q, Yan Z, Chen W, Yan CY, Huang X, Zhang J, Yang PY, Deng HT, Wang JW *et al.* 2012. Structural basis of ultraviolet-B perception by UVR8. *Nature* 484: 214–U296.
- Yang D-L, Yao J, Mei C-S, Tong X-H, Zeng L-J, Li Q, Xiao L-T, Sun T-P, Li J, Deng X-W *et al.* 2012. Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade. *Proceedings of the National Academy of Sciences, USA* 109: E1192–E1200.
- Zavala JA, Mazza CA, Dillon FM, Chludil HD, Ballaré CL. 2014. Soybean resistance to stink bugs (*Nezara viridula* and *Piezodorus guildinii*) increases with exposure to solar UV-B radiation and correlates with isoflavonoid content in pods under field conditions. *Plant, Cell & Environment*. doi: 10.1111/pce.12368.

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Effect of experimental conditions on the response to UV-B radiation for the GO categories discussed in this study.

Fig. S2 Effects of UVR8 in the responses to UV-B reported in Fig. 2.

Table S1 GO categories (<http://geneontology.org/>) used in this study

Table S2 Microarray experiments used in this study

Table S3 Effects of supplemental FR and UV-B on the *Arabidopsis* transcriptome

Methods S1 Additional methods on the physiological experiments and microarray meta-analysis.

Please note: Wiley Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.

Key words: flavonoid, herbivory, jasmonate, meta-analysis, phytochrome, red : far-red ratio, UV-B, UVR8.